



POSTER PRESENTATION

Open Access

The antisense protein of HTLV-2 positively modulates HIV-1 replication

Cynthia Torresilla^{1*}, Sonia Do Carmo¹, Émilie Larocque¹, Estelle Douceron², Jean-Michel Mesnard³, Renaud Mahieux², Benoit Barbeau¹

From 16th International Conference on Human Retroviruses: HTLV and Related Viruses Montreal, Canada. 26-30 June 2013

Unlike HTLV-1, HTLV-2 does not induce leukemia and has been tentatively associated with an HTLV-1-associated myelopathy-like disorder. It has been reported that HTLV-2/HIV-1 co-infected patients progress less rapidly to AIDS than HIV-1-infected individuals. Tax2 has been suggested to mediate this protective state by inducing MIP-1 α expression and blocking HIV-1 infection. As cells from HTLV-2-infected individuals mainly express Antisense Protein 2 (APH-2), our objective was to determine if this protein might also intervene in controlling HIV-1 replication in dually infected individuals. Using Jurkat cells, we first demonstrated that both APH-2 and HBZ, the HTLV-1 analogue, equally induced MIP-1 α in unstimulated and stimulated Jurkat T cells. To assess if APH-2 might directly affect HIV-1 replication, a full length luciferase-expressing proviral DNA was tested in Jurkat cells. Surprisingly, upon co-transfection with an APH-2 expression vector, an increase in luciferase activity was observed, while HBZ expression rather led to reduced reporter gene expression. Western blot analyses and ELISA assay further indicated that HIV-1 p24 levels were more important in APH-2-expressing cells. To determine if APH-2 was directly modulating HIV-1 LTR activity, both NF- κ B and NFAT were tested in stimulated Jurkat cells. Unexpectedly, HBZ and APH-2 inhibited NF- κ B and NFAT activation, albeit at different extent. In addition, LTR activation was also inhibited by both antisense proteins although APH-2 had a more modest effect. Our results thus highlight the complex interplay between HTLV antisense transcript-encoded proteins and HIV-1 expression and further studies will be required to determine the potential impact of APH-2 in HTLV-2/HIV-1-infected individuals.

* Correspondence: cynthia.torresilla@gmail.com

¹Département des sciences biologiques and Centre de recherche BioMed, Université du Québec à Montréal, Montréal (Québec) Canada
Full list of author information is available at the end of the article

Authors' details

¹Département des sciences biologiques and Centre de recherche BioMed, Université du Québec à Montréal, Montréal (Québec) Canada. ²Oncogenèse Rétrovirale, Ligue Nationale Contre Le Cancer, CIRI, INSERM U1111-CNRS UMR5308, Université Lyon 1, Ecole Normale Supérieure de Lyon, LabEx ECOFECT, Lyon, Cedex 07, France. ³Université Montpellier 1, Centre d'études d'agents Pathogènes et Biotechnologies pour la Santé, CNRS, UMS236, Montpellier, France.

Published: 7 January 2014

doi:10.1186/1742-4690-11-S1-P118

Cite this article as: Torresilla et al.: The antisense protein of HTLV-2 positively modulates HIV-1 replication. *Retrovirology* 2014 **11**(Suppl 1): P118.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

