



ORAL PRESENTATION

Open Access

Crossing of the intestinal barrier by HTLV-1 infected lymphocytes

Florent Percher^{1,2,3}, Patricia Jeannin^{1,2}, Pierre-Guillaume Deliège^{1,2}, Antoine Gessain^{1,2}, Aurore Vidy-Roche^{1,2}, Philippe V Afonso^{1,2}, Pierre-Emmanuel Ceccaldi^{1,2,3*}

From 17th International Conference on Human Retroviruses: HTLV and Related Viruses
Trois Ilets, Martinique. 18-21 June 2015

In several areas of high endemicity, HTLV-1 can be transmitted from mother to child, through prolonged breast-feeding. This way of transmission is particularly linked to the development of Adult T Cell Leukemia (ATL). In this context, we studied the mechanisms of HTLV-1 transmission through the digestive tract. We previously demonstrated in an *in vitro* model of compartmentalized co-culture device of human enterocytes, lymphocytes and dendritic cells, that HTLV-1 was unable to infect enterocytes, or disrupt tight cell junctions, but could rather be transported by transcytosis to infect sub-epithelial dendritic cells (Martin-Latil *et al.*, *Blood*, 2012). Since HTLV-1 infection has been shown to be more efficient through cell-associated virions, we also focused on the crossing of HTLV-1 infected lymphocytes through the intestinal barrier. We thus studied the migration of HTLV-1 infected lymphocytes across the intestinal epithelium, both *ex vivo* (on intestinal explant model) and *in vitro* (on a monolayer of human enterocytic cell line cultured on Transwell devices). We demonstrate that, in both systems, infected lymphocytes are able to cross the epithelial barrier, with increased capacities compared to uninfected lymphocytes. Our work suggests that HTLV-1 infected lymphocytes efficiently cross the intestinal barrier, which could constitute an additional way for mother to child HTLV-1 transmission during breast-feeding.

Authors' details

¹Institut Pasteur, Unité d'Epidémiologie et Physiopathologie des Virus Oncogènes, Paris, France. ²CNRS, UMR 3569, Paris, France. ³Université Paris Diderot, Cellule Pasteur, Paris, France.

* Correspondence: pierre-emmanuel.ceccaldi@pasteur.fr

¹Institut Pasteur, Unité d'Epidémiologie et Physiopathologie des Virus Oncogènes, Paris, France

Full list of author information is available at the end of the article

Published: 28 August 2015

doi:10.1186/1742-4690-12-S1-O33

Cite this article as: Percher *et al.*: Crossing of the intestinal barrier by HTLV-1 infected lymphocytes. *Retrovirology* 2015 **12**(Suppl 1):O33.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

