

Poster presentation

Open Access

PI6-24. Crucial contribution of sub-dominant HLA-C allele restricted CTL responses to the control of HIV

J Ibarondo*¹, R Zúñiga², M Farfán², J Suarez², B Mothe¹, A Llano¹, JJ Szinger³, W Hildebrand⁴, J Sánchez⁵, BT Korber³ and C Brander¹

Address: ¹HIVACAT, Fundació irsiCaixa, Badalona, Spain, ²Asociación Civil IMPACTA Salud y Educación, Lima, Peru, ³Los Alamos National Laboratory, Los Alamos, NM, USA, ⁴University of Oklahoma Medical Center, Oklahoma City, OK, USA and ⁵Investigaciones Médicas en Salud (INMENZA), Lima, Peru

* Corresponding author

from AIDS Vaccine 2009
Paris, France. 19–22 October 2009

Published: 22 October 2009

Retrovirology 2009, 6(Suppl 3):P253 doi:10.1186/1742-4690-6-S3-P253

This abstract is available from: <http://www.retrovirology.com/content/6/S3/P253>

© 2009 Ibarondo et al; licensee BioMed Central Ltd.

Background

HLA-A and -B alleles have been associated with relative control of HIV disease and specific HLA-B restricted CTL responses have been implicated with effective antiviral activity. Despite recent reports finding genomic associations between the C locus and relative HIV control, little is known regarding HLA-C restricted CTL responses to HIV and their ability to facilitate viral control.

Methods

In a cohort of 248 HIV infected individuals from Peru, we determined the breadth, magnitude and specificity of dominant and sub-dominant HLA-C restricted immune responses using an overlapping peptide set spanning the entire viral clade B HIV proteome.

Results

After correction for linkage disequilibrium with other HLA class I alleles, HLA-C alleles showed a wide range of relative protection from HIV disease progression, with HLA-C07, C08 and C17 alleles being associated with relative viral control while HLA-C04 was associated with high viral loads and low CD4 counts independently of HLA-B35 co-expression. Assessing "break-through" individuals, i.e. subjects who failed to control in vivo viral replication despite expressing one of the protective HLA-C alleles, indicate that relative control is in at least some cases mediated by sub-dominant T cell responses to HLA-C restricted epitopes.

Conclusion

These analyses strongly suggest that HLA-C restricted immune responses should be considered for HIV vaccine immunogen design, particularly also in the light of recent data indicating potentially synergistic effects of HLA-C restricted epitopes and recognition by NK cells.