

Poster presentation

Open Access

## PI6-53. Distinct subsets of memory T lymphocytes from HIV-1-infected subjects secrete IFN- $\gamma$ and IL-2 in response to novel CD4+ T-cell HIV-1 epitopes

S Fonseca\*<sup>1</sup>, A Coutinho-Silva<sup>1</sup>, D Rodrigues<sup>2</sup>, L Marti<sup>3</sup>, C Moreira-Filho<sup>3</sup>, A Segurado<sup>4</sup>, E Kallás<sup>5</sup>, J Kalil<sup>1</sup> and E Cunha-Neto<sup>1</sup>

Address: <sup>1</sup>Heart Institute, University of São Paulo, São Paulo, Brazil, <sup>2</sup>Instituto Clemente Ferreira, São Paulo-SP, Brazil, <sup>3</sup>Programa Interunidades em Biotecnologia USP-IPT-Butantan, São Paulo-SP, Brazil, <sup>4</sup>Divisão de Doenças Infecciosas, FMUSP, São Paulo, Brazil and <sup>5</sup>Divisão de Imunologia Clínica e Alergia da FMUSP, São Paulo, Brazil

\* Corresponding author

from AIDS Vaccine 2009  
Paris, France. 19–22 October 2009

Published: 22 October 2009

Retrovirology 2009, 6(Suppl 3):P282 doi:10.1186/1742-4690-6-S3-P282

This abstract is available from: <http://www.retrovirology.com/content/6/S3/P282>

© 2009 Fonseca et al; licensee BioMed Central Ltd.

### Background

Functional memory T lymphocytes are involved in the control of HIV replication. Recently, we have described 18 HLA-DR promiscuous epitopes from HIV-1 consensus B sequences. Some of them contain CD8 epitopes inserted.

### Objective

To investigate the functionality of the subsets of memory T cells that respond to these peptides among PBMC from HIV+ patients.

### Methods

PBMC from 35 HIV+ patients with CD4 count >250 cells/ $\mu$ L: 8 aviremics in ART (AV-ART), 8 viremics in ART (VI-ART) (VL = 2260–31000 copies/mL) and 9 viremics naïve of therapy (VI-NO ART) (VL = 17100–698000 copies/mL) and 7 healthy donors were stimulated with pooled HIV- or CMV-peptides. The cellular phenotype and cytokine production were evaluated by multiparameter flow cytometry.

### Results

The phenotypic analysis showed a significant reduction of CD8+ T naïve cells in VI-ART ( $P = 0.014$ ) and VI-NO ART ( $P = 0.003$ ) compared to healthy group. The functional analysis revealed that all HIV-infected patients of our study had CD4+ and CD8+ secreting IFN- $\gamma$  and/or IL-2 in

response to our pool of HIV-1 peptides. This recognition involved central memory (TCM, CCR7+CD45RA-), effector memory (TEM, CCR7-CD45RA-), and CD45RA+ effector memory (TEMRA, CCR7-CD45RA+). In the CD4+ compartment the VI-ART group showed the smallest frequency of functional memory in response to our HIV pooled peptides. This observation was more evident in respect to IL-2 producing cells.

### Conclusion

Our results suggest a functional heterogeneity of T memory subsets in response to set HIV-1 peptides, which may constitute a new candidate to an anti-HIV vaccine.