

Poster presentation

Open Access

PI7-22. Impact of rare adenovirus seroprevalence on HIV-1 acquisition in the Step study

LF Maxfield*¹, SL King¹, AM Riggs¹, R Dilan¹, A LaPorte¹, J Hural², MJ McElrath², J Goudsmit³ and DH Barouch¹

Address: ¹Division of Viral Pathogenesis, Beth Israel Deaconess Medical Center, Boston, MA, USA, ²HVTN, Vaccine & Infectious Disease Inst, Fred Hutchinson Cancer Res Ctr, Seattle, WA, USA and ³Crucell, Leiden, Netherlands

* Corresponding author

from AIDS Vaccine 2009
Paris, France. 19–22 October 2009

Published: 22 October 2009

Retrovirology 2009, 6(Suppl 3):P304 doi:10.1186/1742-4690-6-S3-P304

This abstract is available from: <http://www.retrovirology.com/content/6/S3/P304>

© 2009 Maxfield et al; licensee BioMed Central Ltd.

Background

In the phase IIb Step study, vaccinees with baseline Ad5-specific neutralizing antibodies (NABs) exhibited a potential increased rate of HIV-1 acquisition as compared with placebo controls. We sought to evaluate whether baseline Ad5 NABs correlated with NABs to non-type C rare Ad serotypes and whether baseline NABs to rare Ad serotypes correlated with enhancement of HIV-1 acquisition after rAd5-gag/pol/nef vaccination.

Methods

In a case-controlled study, baseline sera from 81 cases who acquired HIV-1 infection during the Step study (52 vaccinees and 29 placebo recipients) and 324 non-cases who did not acquire HIV-1 infection (208 vaccinees and 29 placebo recipients) were evaluated for Ad5, Ad26, Ad35, and Ad48 NAB titers. Non-cases were randomly selected and matched for treatment arm, Ad5 serostatus, circumcision status, and geographic region.

Results

Baseline Ad5 seroprevalence was 58% in these samples with a median titer of 702. In contrast, baseline Ad26, Ad35, and Ad48 seroprevalence was 14%, 6%, and 17%, respectively, with median titers of 49, 47, and 46. There were no correlations between baseline Ad5 NABs and Ad26/35/48 NABs ($P = 0.44-0.77$, Spearman rank correlation tests). Moreover, there were no differences observed

in baseline Ad26/35/48 NABs between cases and non-cases ($P = 0.46-0.97$, Wilcoxon rank sum tests).

Conclusion

Ad5 NABs and Ad26/35/48 NABs were independent variables that did not co-segregate, indicating that Ad5 NABs were not a surrogate marker for and did not predict NABs to these rare Ad serotypes. In addition, no increased HIV-1 acquisition was observed in subjects with baseline Ad26/35/48 NABs following rAd5 vaccination. These data suggest that cross-reactive immunity against heterologous Ad serotypes did not contribute to increased HIV-1 susceptibility in this study. These findings have important implications for the development of rare serotype Ad vector-based vaccines for HIV-1.

Acknowledgements

NIAID/HVTN, NIAID/IPCAVD, Gates/CAVD.