



POSTER PRESENTATION

Open Access

# The immunoglobulin CH1 constant region modulates antigen binding affinity and functional activities of the broadly neutralizing 2F5 HIV specific antibody

Daniela Tudor<sup>1\*</sup>, Anne-Sophie Drillet<sup>1</sup>, Isabelle Schwartz-Cornil<sup>2</sup>, Ruizhong Shen<sup>3</sup>, Phillip D Smith<sup>3</sup>, Morgane Bomsel<sup>1</sup>

From 16<sup>th</sup> International Symposium on HIV and Emerging Infectious Diseases  
Marseille, France. 24-26 March 2010

## Background

The ability of the heavy chain constant region (CH) to affect antibody affinity and specificity could be at the origin of a stronger or weaker memory response, depending on the isotype. Using as a model the broadly neutralizing human mAb 2F5, directed against the membrane proximal region (MPER) of the HIV-1 envelope transmembrane subunit gp41, we investigated the interplay between 2F5 isotype and functional activity.

## Methods

A 2F5 IgA isotype was constructed from the 2F5 IgG1. Functional monomeric 2F5 IgA and IgG1 were expressed in CHO cells and their immunochemical characteristics and anti-HIV-1 in-vitro activity were evaluated.

## Results

As compared to 2F5 IgG1, 2F5 IgA sharing identical VH and VL domains but in a different CH context: (i) binds with higher affinities gp41 and MPER peptides; (ii) has an increased capacity at inhibiting endocytosis of HIV-1 by dendritic cells; (iii) has an increased HIV-1 neutralizing activity in lymphocytic CD4+ T cells; (iv) blocks more efficiently HIV-1 transcytosis across epithelial monolayers in-vitro and normal human rectal mucosa, but (v) has lower ADCC activity. Epitope mapping with a 7 mer epitope library shows that 2F5 IgA recognizes

essentially the same hexapeptide epitope as its IgG counterpart.

## Discussion

These results show that the CH region can fine-tune the specificity of an antibody, by modulating its binding affinity to the antigen and the neutralizing activity of variable-region of otherwise identical antibodies. The determinant role of CH region on affinity and specificity changes our understanding of vaccine responses. In the context of HIV-1, which is mainly transmitted sexually, these results strongly suggest that raising a mucosal humoral IgA based response will be superior to an IgG one in blocking HIV-1 transmission.

## Author details

<sup>1</sup>Institut Cochin, Paris, France. <sup>2</sup>INRA, Jouy-en-Josas, France. <sup>3</sup>University of Birmingham, Alabama, USA.

Published: 11 May 2010

doi:10.1186/1742-4690-7-S1-P23

**Cite this article as:** Tudor *et al.*: The immunoglobulin CH1 constant region modulates antigen binding affinity and functional activities of the broadly neutralizing 2F5 HIV specific antibody. *Retrovirology* 2010 **7**(Suppl 1):P23.

\* Correspondence: [daniela.tudor@inserm.fr](mailto:daniela.tudor@inserm.fr)

<sup>1</sup>Institut Cochin, Paris, France