



MEETING ABSTRACT

Open Access

# Localization and sub-cellular shuttling of HTLV-1 Tax with the RNAi machinery component Droscha

Rachel Van Duyne<sup>1,2</sup>, Irene Guendel<sup>1</sup>, Mudit Tyagi<sup>1</sup>, Aarthi Narayanan<sup>1</sup>, Zachary Klase<sup>3</sup>, Kylene Kehn-Hall<sup>1</sup>, John Semmes<sup>4</sup>, Fatah Kashanchi<sup>1,2\*</sup>

From 15th International Conference on Human Retroviruses: HTLV and Related Viruses  
Leuven and Gembloux, Belgium. 5-8 June 2011

The innate ability of the human cell to silence endogenous retroviruses through RNA sequences encoding micro-RNAs, suggests that the cellular RNAi machinery is a major mean by which the host mounts a response against contemporary retroviruses, such as HIV-1 and HTLV-1. Several recent publications have identified cellular miRNAs that target and hybridize to specific sequences of both the HIV-1 and HTLV-1 transcripts. However, much like the variety of host immune responses to retroviral infection, the virus itself contains mechanisms that assist in the evasion of viral inhibition through manipulation of the cellular RNAi pathway. Retroviruses can hijack both the enzymatic and catalytic components of the RNAi pathway, in some cases to produce novel viral miRNAs that can either assist in active viral infection or promote a latent state of infection. Here, we propose that HTLV-1 viral proteins contribute to the dysregulation of the RNAi pathway by altering expression of key components of the pathway. A survey of uninfected and infected cell lines revealed that Droscha was present at lower levels in all HTLV infected lines. Additionally, transfection of HeLa cells with Tax shows colocalization of Tax and Droscha in the nucleus (speckles), suggesting that the HTLV-1 viral transactivator physically interacts with Droscha and targets it to specific areas of the cell. This data suggests the direct interaction of HTLV-1 viral components with RNAi machinery proteins which may lead to their dysregulation in infected cells.

#### Author details

<sup>1</sup>George Mason University, Department of Molecular and Microbiology, National Center for Biodefense and Infectious Diseases, Manassas, VA, 20110,

\* Correspondence: [fkashanc@gmu.edu](mailto:fkashanc@gmu.edu)

<sup>1</sup>George Mason University, Department of Molecular and Microbiology, National Center for Biodefense and Infectious Diseases, Manassas, VA, 20110, USA

Full list of author information is available at the end of the article

USA. <sup>2</sup>The George Washington University Medical Center, Department of Microbiology, Immunology, and Tropical Medicine, Washington, DC, 20037, USA. <sup>3</sup>National Institutes of Health, Molecular Virology Section, Laboratory of Molecular Microbiology, NIAID, Bethesda, Maryland, 20892, USA. <sup>4</sup>Eastern Virginia Medical School, Department of Microbiology and Molecular Cell Biology, Cancer Biology and Infectious Disease Research Center, Norfolk, VA, 23508, USA.

Published: 6 June 2011

doi:10.1186/1742-4690-8-S1-A159

Cite this article as: Van Duyne *et al.*: Localization and sub-cellular shuttling of HTLV-1 Tax with the RNAi machinery component Droscha. *Retrovirology* 2011 **8**(Suppl 1):A159.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

