



MEETING ABSTRACT

Open Access

HTLV-I proviral load in Argentinean subjects with indeterminate western blot patterns

Andrea Mangano^{1*}, Natalia Altamirano¹, Mirta Remesar², María B Bouzas³, Paula Aulicino¹, Inés Zapiola³, Ana D Pozo², Luisa Sen¹

From 15th International Conference on Human Retroviruses: HTLV and Related Viruses Leuven and Gembloux, Belgium. 5-8 June 2011

Background

A considerable high proportion of HTLV-I/II seroindeterminate blood donors have been documented in many countries including Argentina. In 5-10% of indeterminate Western Blot (WB) cases proviral sequences are detected. The aim of the study was to evaluate HTLV-I proviral load (pVL) in WB indeterminate PCR positive cases.

Methodology

A total of 87 indeterminate WB samples (HTLV blot 2.4 assay- Genelabs Diagnostics-) were studied over a 10 year period referred from the Blood Bank at Garrahan Hospital (n=83) and from the Virology Unit at Muñiz Hospital (n=4). HTLV-I and -II tax and pol proviral sequences were amplified by in-house nested PCR assays. HTLV-I pVL was estimated by a quantitative real-time PCR assay targeting the HTLV-I pol gene and the albumin gene as normalizer. The limit of detection of the assay was 2.6 log₁₀ copies of HTLV-I/106 PBMCs (0.04% copies/100 PBMCs).

Results

In 8/87 samples HTLV proviral sequences were amplified, 7 HTLV-I and one HTLV-II. Proviral load was detectable in 4 of the 7 HTLV-I positive samples ranging from 2.72 log₁₀ copies/106 PBMCs (0.05%) to 4.65 log₁₀ copies/106 PBMCs (4.5%), but was undetectable in the remaining 3 (<0.04%). One of the subjects followed for 5 years remained with low pVL (median = 0.36%, range 0.05-1.43%) and without changes in the WB profile.

Conclusions

Detectable HTLV-I proviral load in half of the cases with indeterminate WB profiles with positive PCR underlies the importance to conduct follow-up studies to evaluate the evolution of the infection.

Author details

¹Laboratorio de Biología Celular y Retrovirus-CONICET, Hospital "J. P. Garrahan", Ciudad Autónoma de Buenos Aires, Argentina. ²Servicio de Hemoterapia, Hospital "J. P. Garrahan", Ciudad Autónoma de Buenos Aires, Argentina. ³Unidad de Virología, Hospital Muñiz, Ciudad Autónoma de Buenos Aires, Argentina.

Published: 6 June 2011

doi:10.1186/1742-4690-8-S1-A245

Cite this article as: Mangano *et al.*: HTLV-I proviral load in Argentinean subjects with indeterminate western blot patterns. *Retrovirology* 2011 8 (Suppl 1):A245.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: amangano@garrahan.gov.ar

¹Laboratorio de Biología Celular y Retrovirus-CONICET, Hospital "J. P. Garrahan", Ciudad Autónoma de Buenos Aires, Argentina
Full list of author information is available at the end of the article