



MEETING ABSTRACT

Open Access

# Evidence of STLV-2 and STLV-3 infections in wild-living bonobos (*P. paniscus*) from the Democratic Republic of Congo

Steve Ahuka-Mundeke<sup>1,2</sup>, Florian Lieg ois<sup>1</sup>, Octavie Lunguya<sup>2</sup>, Valentin Mbenzo<sup>3</sup>, Menard Mbende<sup>3</sup>, Bila-Isia Inogwabini<sup>3</sup>, Jean-Jacques Muyembe<sup>2</sup>, Eric Delaporte<sup>1</sup>, Martine Peeters<sup>1\*</sup>

From 15th International Conference on Human Retroviruses: HTLV and Related Viruses  
Leuven and Gembloix, Belgium. 5-8 June 2011

## Background

Among the four types of HTLV (1 to 4), only three have their simian counterparts (STLV-1, 2 and 3). STLV-1 and 3 have been found in a large number of captive and wild-living monkeys and great apes from Africa and Asia. STLV-2 was reported only in a limited number of captive bonobos (*P. paniscus*) from Democratic Republic of Congo (DRC) and despite its genetic distance with HTLV-2, STLV-2 is considered as a simian counterpart of HTLV-2. To date, no evidence of STLV-2 or any other known STLVs have been documented yet in wild-living bonobos. Given that bonobos are an endangered species, investigation of pathogens in this species in the wild is only possible by non-invasive sampling. Here we report a survey aimed at characterising simian retroviruses from wild-living bonobos in DRC.

## Methods

Between March and July 2010, fecal samples from wild-living bonobos (*P. paniscus*) were collected at Malebo forest in the Bandundu province located in the western part of DRC. Samples were collected in RNA-later. Species confirmation was done by mtDNA analyses. All faecal samples were screened for STLV infection using a generic PCR allowing amplification of partial fragment in tax (220bp). Sequence and phylogenetic analysis was done to confirm the STLV infection and identify STLV types. Attempts to amplify other gene fragments were done to further characterize the new STLV viruses.

## Results

A total of 268 fecal samples were collected in 2010 and mitochondrial DNA analyses confirmed that all the samples were from bonobos (*P. paniscus*). Overall, 3 (1.1%) out of 268 samples yielded positive amplification of tax fragment. Among them, one (Pp5538) was identified as STLV-2 and two (Pp5489 and Pp5560) as STLV-3 by phylogenetic analysis of tax fragment. Phylogenetic analyses of tax and LTR fragments showed that the new STLV-2 from Pp5538 clustered with STLV-2 strains previously described in captive bonobos.

All PCR attempts to amplify the LTR fragment of Pp5489 and 5560 samples were unsuccessful.

## Conclusion

Our study shows that faecal samples can be used to screen for STLV infection in endangered apes, although most likely with lower sensitivities. We confirmed STLV-2 infection in wild-living bonobos and showed for the first time that bonobos are also infected with STLV-3. Additional work is warranted to fully characterize the new STLV-2 and 3 strains and to document the STLV diversity on a larger number of wild bonobos from different geographic regions.

## Author details

<sup>1</sup>UMI 233, Institut de Recherche pour le Developpement and University of Montpellier 1, Montpellier, France. <sup>2</sup>National Institute of Biomedical Research, Kinshasa, Democratic Republic of Congo. <sup>3</sup>Lac Tumba Project, WWF, Kinshasa, Democratic Republic of Congo.

<sup>1</sup>UMI 233, Institut de Recherche pour le Developpement and University of Montpellier 1, Montpellier, France

Full list of author information is available at the end of the article

Published: 6 June 2011

doi:10.1186/1742-4690-8-S1-A92

**Cite this article as:** Ahuka-Mundeke *et al.*: Evidence of STLV-2 and STLV-3 infections in wild-living bonobos (*P. paniscus*) from the Democratic Republic of Congo. *Retrovirology* 2011 **8**(Suppl 1):A92.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

