



POSTER PRESENTATION

Open Access

# The dual (activating/suppressive) effect of extracellular TatHIV-1 is driven by the inflammatory microenvironment of infected lymphoid foci

Hélène Le Buanec<sup>1,2,3</sup>, Thomas Sené<sup>1,2,3</sup>, Armand Bensussan<sup>1,2,3</sup>, Robert Gallo<sup>4</sup>, Daniel Zagury<sup>5\*</sup>

From 17th International Symposium on HIV and Emerging Infectious Diseases (ISHEID) Marseille, France. 23-25 May 2012

It has been shown that HIV-1 infects activated but not resting CD4+ T cells [1] and that CPE induced by viral replication together with the immunosuppressive effect triggered by extracellular Tat protein [2] account for the decrease of CD4+ T cell count in infected patients. In lymphoid foci, dependent on the level of viral infection, the stromal microenvironment surrounding immune cells could include, together with extracellular Tat [3] and circulating antiviral IFN- $\alpha$ , inflammatory innate factors such as ATP and derivatives released by CPE-derived dead cells.

We show that, according to its concentration and the presence of inflammatory factors (IFN- $\alpha$ , ATP and ATP-derivatives), Tat protein may exert either an activation with enhanced production of IL2 or an immune suppression of stimulated CD4+ T cells subpopulations.

The double-edged sword of Tat activity on CD4+ T cells could account for its immunopathogenic effects both at the early stage of infection (by allowing CD4+ T cells activation and viral replication) and at late stages (by inducing immunosuppression, source of opportunistic infections). Indications for targeting Tat protein by therapeutic vaccines in subgroups of HIV-1 infected patients will be discussed.

#### Author details

<sup>1</sup>INSERM U976, F-75475, Paris, France. <sup>2</sup>Université Paris Diderot, Sorbonne Paris Cité, Laboratory of Immunology, Dermatology & Oncology, UMR-S 976, F-75475, Paris, France. <sup>3</sup>Service de dermatologie, Hopital Saint Louis, F-75010,

Paris, France. <sup>4</sup>Institute of Human Virology, University of Maryland Baltimore, Maryland, USA. <sup>5</sup>Neovacs SA, Paris, France.

Published: 25 May 2012

#### References

1. Zagury D, *et al*: Long-term cultures of HTLV-III-infected T cells: a model of cytopathology of T-cell depletion in AIDS. *Science* 1986, **231**(4740):85.
2. Viscidi RP, *et al*: Inhibition of antigen-induced lymphocyte proliferation by Tat protein from HIV- Science. 1989, **246**(4937):1606.
3. Ensoli B, *et al*: Release, uptake, and effects of extracellular human immunodeficiency virus type 1 Tat protein on cell growth and viral transactivation. *J Virol* 1993, **67**(1):277.

doi:10.1186/1742-4690-9-S1-P35

Cite this article as: Le Buanec *et al*: The dual (activating/suppressive) effect of extracellular TatHIV-1 is driven by the inflammatory microenvironment of infected lymphoid foci. *Retrovirology* 2012 **9**(Suppl 1):P35.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit



\* Correspondence: dzagury@neovacs.com

<sup>5</sup>Neovacs SA, Paris, France

Full list of author information is available at the end of the article