



POSTER PRESENTATION

Open Access

# Characterisation of a spumavirus Gag protein

David C Goldstone<sup>1</sup>, Thomas G Flower<sup>1</sup>, Neil J Ball<sup>1\*</sup>, Marta Sanz-Ramos<sup>2</sup>, Melvin W Yap<sup>2</sup>, Roksana W Ogradowicz<sup>1</sup>, Nicole Stanke<sup>3</sup>, Juliane Reh<sup>3</sup>, Dirk Lindemann<sup>3</sup>, Jonathan P Stoye<sup>2</sup>

From *Frontiers of Retrovirology: Complex retroviruses, retroelements and their hosts*  
Cambridge, UK. 16-18 September 2013

## Background

Human prototypic foamy virus (HPFV) belongs to the *spumaretrovirinae* subfamily and is an attractive vector candidate for gene therapy [1] as it is apathogenic. The Gag protein is not cleaved into matrix (MA), capsid (CA) and nucleocapsid (NC) as occurs in orthoretroviruses; rather, it is able to perform the roles of these proteins as a single polypeptide [2]. Foamy virus Gag proteins are targets for restriction factors such as Trim5 $\alpha$  [3] and also interact with the aminoterminal leader peptide of the envelope protein (Env). This Gag-Env interaction is essential for budding of viral particles from the host cell [4,5].

## Materials and methods

To investigate HPFV Gag assembly and Gag-Env interactions we have undertaken combined structural and biophysical studies using X-ray crystallography, NMR, multi-angle laser light scattering (MALLS) and analytical ultracentrifugation (AUC).

## Results

An N-terminal domain (NtD) of HPFV-Gag has been identified which is functionally related to CA and MA in other retroviruses despite very low sequence homology. Structural and solution studies of the HPFVGagNtD reveal that it forms a very stable homodimer which interacts through an extended coiled-coil region. A separate region of HPFV-Gag that exhibits a propensity for self-association has been identified and preliminary structural and biophysical analysis suggests that it is able to form homodimers with a dissociation constant of approximately 30  $\mu$ M. The molecular basis of the Gag-Env interaction has been structurally characterised and mutagenic studies, both *in vitro* and *in vivo*, have revealed the

importance of the individual residues involved in this interaction.

## Conclusions

In stark contrast to orthoretroviral capsid which self-associates to form a predominantly hexameric lattice, the HPFV-Gag-NtD forms stable homodimers in solution. The identification of a secondary self-association region suggests that these dimers may undergo further multimerisation in the full-length Gag protein. Comparison of Trim5 $\alpha$  specificity between foamy viruses suggests that the recognition site resides on the exposed surface of the N-terminal domain of the Gag protein.

## Authors' details

<sup>1</sup>Division of Molecular Structure, MRC National Institute for Medical Research, The Ridgeway, Mill Hill, London, NW7 1AA, UK. <sup>2</sup>Division of Virology, The Ridgeway Mill Hill, London, NW7 1AA, UK. <sup>3</sup>Institute of Virology, Technische Universität Dresden, Fetscherstraße 74, 01307 Dresden, UK.

Published: 19 September 2013

## References

1. Lindemann D, Rethwilm A: **Foamy virus biology and its application for vector development.** *Viruses* 2011, **3**:561-585.
2. Flügel R, Pfrepper K: **Proteolytic processing of foamy virus Gag and Pol proteins.** *Curr Top Microbiol Immunol* 2003, **277**:63-88.
3. Yap M, Lindemann D, Stanke N, Reh J, Westphal D, Hanenberg H, Ohkura S, Stoye J: **Restriction of foamy viruses by primate Trim5 $\alpha$ .** *J Virol* 2008, **82**:5429-5439.
4. Wilk T, Geiselhart V, Frech M, Fuller S, Flügel R, Löchelt M: **Specific interaction of a novel foamy virus Env leader protein with the N-terminal Gag domain.** *J Virol* 2001, **75**:7995-8007.
5. Pietschmann T, Heinkelein M, Heldmann M, Zentgraf H, Rethwilm A, Lindemann D: **Foamy virus capsids require the cognate envelope protein for particle export.** *J Virol* 1999, **73**:2613-2621.

doi:10.1186/1742-4690-10-S1-P3

Cite this article as: Goldstone et al.: Characterisation of a spumavirus Gag protein. *Retrovirology* 2013 **10**(Suppl 1):P3.

<sup>1</sup>Division of Molecular Structure, MRC National Institute for Medical Research, The Ridgeway, Mill Hill, London, NW7 1AA, UK  
Full list of author information is available at the end of the article