



POSTER PRESENTATION

Open Access

Extreme genetic fragility of the HIV-1 capsid

Suzannah Rihn^{1,2*}, Sam Wilson^{1,5}, Nick Loman⁴, Mudathir Alim^{1,2}, Saskia Bakker⁵, David Bhella⁵, Frazer Rixon⁵, Paul Bieniasz^{1,3}

From *Frontiers of Retrovirology: Complex retroviruses, retroelements and their hosts*
Cambridge, UK. 16-18 September 2013

Genetic robustness, or fragility, is defined as the ability, or lack thereof, of a biological entity to maintain function in the face of mutations. Viruses that replicate via RNA intermediates exhibit high mutation rates, and robustness should be particularly advantageous to them. The capsid (CA) domain of the HIV-1 Gag protein is under strong pressure to conserve functional roles in viral assembly, maturation, uncoating, and nuclear import. However, CA is also under strong immunological pressure to diversify in hosts. Therefore, it would be particularly advantageous for CA to evolve genetic robustness. To probe the genetic robustness of HIV-1 CA, we generated a library of single amino acid substitution mutants, encompassing almost half the residues in CA. Strikingly, we found HIV-1 CA to be the most genetically fragile protein that has been analyzed using such an approach, with 70% of mutations yielding replication-defective viruses. Although CA participates in several steps in HIV-1 replication, analysis of conditionally (temperature sensitive) and constitutively non-viable mutants revealed that the biological basis for its genetic fragility was primarily the need to coordinate the efficient and proper assembly of a mature viral capsid. All mutations that exist in naturally occurring HIV-1 subtype B populations at a frequency >3%, and were also present in the mutant library, had fitness levels that were >40% of WT. However, some mutations with high fitness did not occur in natural populations, suggesting another form of selection pressure limiting variation *in vivo*. Additionally, known protective CTL epitopes occurred preferentially in domains of the HIV-1 CA that were even more genetically fragile than HIV-1 CA as a whole. The extreme genetic fragility of HIV-1 CA may be one reason why cell-mediated immune responses to Gag correlate with better prognosis in HIV-1 infection, and suggests that CA is a good target for therapy and vaccination strategies.

Authors' details

¹Laboratory of Retrovirology, The Rockefeller University, New York, NY, USA. ²Aaron Diamond AIDS Research Center, The Rockefeller University, New York, NY, USA. ³Howard Hughes Medical Institute, Aaron Diamond AIDS Research Center, New York, NY, USA. ⁴Centre for Systems Biology, University of Birmingham, Birmingham, UK. ⁵MRC Centre for Virus Research, Institute of Infection, Immunity and Inflammation, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow, UK.

Published: 19 September 2013

doi:10.1186/1742-4690-10-S1-P73

Cite this article as: Rihn et al.: Extreme genetic fragility of the HIV-1 capsid. *Retrovirology* 2013 **10**(Suppl 1):P73.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Laboratory of Retrovirology, The Rockefeller University, New York, NY, USA
Full list of author information is available at the end of the article