



POSTER PRESENTATION

Open Access

Naturally occurring protease inhibitor resistance mutations and their frequencies in HIV proviral sequences of drug-naïve sex workers in Nairobi, Kenya

Raghavan Sampathkumar^{1,3}, Elnaz Shadabi^{1,3}, David La³, John Ho³, Binhua Liang^{1,3}, Joshua Kimani², Terry B Ball^{1,3}, Francis A Plummer^{1,3}, Ma Luo^{1,3*}

From 16th International Conference on Human Retroviruses: HTLV and Related Viruses Montreal, Canada. 26-30 June 2013

Sub-Saharan Africa accounts for 69% of the people living with HIV globally. An estimated 1,600,000 Kenyans are living with HIV-1. Antiretroviral therapy (ART) has saved 9 million life-years in Sub-Saharan Africa. However, drug resistance mutations reduce the effectiveness of ART, and need to be monitored for effective ART. Naturally occurring primary antiretroviral drug resistance mutations have not been well analyzed in ART naïve HIV+ patients in Kenya. Here we have examined protease inhibitor (PI) resistance mutations in ART naïve HIV-1 seropositive women in Pumwani sex worker cohort established in Nairobi, Kenya, wherein HIV-1 infection is predominantly caused by subtypes A and D viruses. We have analyzed consensus sequences of HIV protease from 109 drug-naïve patients, as a part of HIV-1 whole-genome sequencing using 454 sequencing methodology. Analysis using HIVdb program revealed a prevalence of 22% (24/109) PI resistance mutations among the study subjects. D30N (3.7%), M46I (0.9%) and V82F (0.9%) are the major mutations observed. D30N mutation is known to confer high-level resistance to nelfinavir. M46I and V82F confer resistance to indinavir, lopinavir, fosamprenavir and nelfinavir. In addition, many minor mutations were found at seven different drug resistance sites. It is important to study the implications of these mutations to the effectiveness of specific PI drug treatment. This study provides valuable data pertaining to primary drug

resistance in Kenyan HIV-1 infected patients before ART became available.

Authors' details

¹Department of Medical Microbiology, University of Manitoba, Winnipeg, Manitoba, Canada. ²Department of Medical Microbiology, University of Nairobi, Nairobi, Kenya. ³National Microbiology Laboratory, Public Health Agency of Canada, Winnipeg, Manitoba, Canada.

Published: 7 January 2014

doi:10.1186/1742-4690-11-S1-P133

Cite this article as: Sampathkumar *et al.*: Naturally occurring protease inhibitor resistance mutations and their frequencies in HIV proviral sequences of drug-naïve sex workers in Nairobi, Kenya. *Retrovirology* 2014 **11**(Suppl 1):P133.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: Ma.Luo@phac-aspc.gc.ca

¹Department of Medical Microbiology, University of Manitoba, Winnipeg, Manitoba, Canada

Full list of author information is available at the end of the article

