



POSTER PRESENTATION

Open Access

BNZ-gamma peptide, a potential therapeutic agent in HTLV-1 associated myelopathy

Raya Massoud¹, Yoshimi Enose-Akahata¹, Yutaka Tagaya², Nazli Azimi³, Steven Jacobson^{1*}

From 16th International Conference on Human Retroviruses: HTLV and Related Viruses Montreal, Canada. 26-30 June 2013

The precise mechanism of HAM/TSP pathogenesis remains unclear however many findings suggest that HTLV-1 can activate the infected lymphocytes in the peripheral circulation leading to enhanced migration into the CNS. Once in this compartment, there is preferential expansion of the infected cells and a compartmentalized interaction between virus-specific CD8+ T-cells and virus- infected CD4+ lymphocytes leading to bystander damage of neural tissues. Dysregulation of T cells activation has been reported in HAM/TSP, particularly dependent on the induction of the stimulatory cytokine loops IL-2/IL-2R α , IL-15/IL-15R α and IL-9/IL-9R γ by the virus. All these cytokines share a common chain receptor, the gamma chain (γ C), offering a potential therapeutic target. BNZ- γ is a 19-mer peptide, that was designed based on the presence of a moderately conserved region shared by all six members of the Gamma cytokine family (IL-2, IL-4, IL-7, IL-9, IL-15 and IL-21) in their D-helix structure. *In vitro*, we showed that BNZ- γ selectively blocks binding and downstream signaling of IL-2, IL-9 and IL-15. In PBMC of HAM/TSP patients, BNZ- γ suppressed *ex-vivo* spontaneous proliferation (SP) in the majority of the tested samples. Additionally a pegylated form of the peptide (N-PEG40) could suppress SP, CD25 expression and STAT5 phosphorylation in the CD8+ T-cells. More importantly, N-PEG40 decreased proliferation of Tax-specific CD8+ cells, a known pathogenic effector cell in HAM/TSP. We are currently studying its effects on functional markers of effector /memory T-cells. These results suggest that manipulation of the inflammatory cytokine loops may be of therapeutic value in the treatment of HAM/TSP.

Authors' details

¹Viral Immunology Section, Neuroimmunology Branch, National Institutes of Neurological Disorders and Stroke, Bethesda, MD, USA. ²Institute of Human Virology, University of Maryland School of Medicine, Baltimore, MD, USA. ³BIONIZ, CA, USA.

Published: 7 January 2014

doi:10.1186/1742-4690-11-S1-P14

Cite this article as: Massoud *et al.*: BNZ-gamma peptide, a potential therapeutic agent in HTLV-1 associated myelopathy. *Retrovirology* 2014 11(Suppl 1):P14.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: jacobsons@ninds.nih.gov

¹Viral Immunology Section, Neuroimmunology Branch, National Institutes of Neurological Disorders and Stroke, Bethesda, MD, USA

Full list of author information is available at the end of the article

