



POSTER PRESENTATION

Open Access

HTLV-1 molecular epidemiology in central Australia: Two distinctive HTLV-1 Subtype C lineages in Indigenous Australians

Olivier Cassar^{1,2*}, Lloyd Einsiedel³, Philippe V Afonso^{1,2}, Antoine Gessain^{1,2}

From 16th International Conference on Human Retroviruses: HTLV and Related Viruses
Montreal, Canada. 26-30 June 2013

HTLV-1 infects approximately 5-10 million people worldwide. It is widely distributed, with clusters of high endemicity in certain geographic areas or ethnic groups. Seven main HTLV-1 subtypes have been described and the Melanesian/Australian subtype C is only found in Oceania. In Australia, HTLV-1 is endemic with ATLL, TSP/HAM, bronchiectasis, ID and severe scabies cases reported. However, HTLV-1 clinical significance is debated and only one partial HTLV-1 sequence from an Aboriginal Australian (MSHR-1) is available. Therefore, establishing a large HTLV-1 sequence database is essential for the understanding of the epidemiology and pathogenesis of this virus in Indigenous people. Samples were obtained from 23 HTLV-1 patients originated from four broad geographical and language-based areas. Sequence comparison indicated that the 23 new HTLV-1 LTR and Tax-Gag sequences belong to the subtype C as MSHR-1 strain. Phylogenetic analyses evidenced two main clades (Solomon Islands/Vanuatu *versus* Australia) within subtype C. Interestingly, the Australian clade can be itself divided into two clusters: the first comprising strains characterized among most of the Indigenous Australians from the North and the second including a majority of individuals originating from the South and Central Australia. Thus, there is a clear evidence of a specific HTLV-1 clustering according to the childhood residence of the HTLV-1 infected individuals, and subsequently their language group affinity. Taking into account that non-human primates have never been found in Australia, we believe that these HTLV-1 subtype C variants have

been present among the first ancestors of modern Indigenous Australians for several tens of millennia.

Authors' details

¹Institut Pasteur, Unité d'Epidémiologie et Physiopathologie des Virus Oncogènes, Département de Virologie, Paris, France. ²CNRS, UMR 3569, Paris, France. ³Flinders University/Northern Territory Rural Clinical School, Alice Springs Hospital, Northern Territory, Australia.

Published: 7 January 2014

doi:10.1186/1742-4690-11-S1-P56

Cite this article as: Cassar *et al.*: HTLV-1 molecular epidemiology in central Australia: Two distinctive HTLV-1 Subtype C lineages in Indigenous Australians. *Retrovirology* 2014 **11**(Suppl 1):P56.

* Correspondence: olivier.cassar@pasteur.fr

¹Institut Pasteur, Unité d'Epidémiologie et Physiopathologie des Virus Oncogènes, Département de Virologie, Paris, France
Full list of author information is available at the end of the article

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

