

Commentary

Open Access

## HTLV-I Tax: centrosome amplification and cancer

Anne Pumfery<sup>1</sup>, Cynthia de la Fuente<sup>2</sup> and Fatah Kashanchi<sup>\*3,4</sup>

Address: <sup>1</sup>Seton Hall University, Department of Biology, South Orange, NJ 07079, USA, <sup>2</sup>The Rockefeller University, Laboratory of Virology and Infectious Disease, New York, NY 10021, USA, <sup>3</sup>The George Washington University Medical Center, Department of Biochemistry and Molecular Biology, Washington, DC 20037, USA and <sup>4</sup>The Institute for Genomic Research, Rockville, Maryland 20850, USA

Email: Anne Pumfery - bcmamp@gwumc.edu; Cynthia de la Fuente - cdelafuent@mail.rockefeller.edu; Fatah Kashanchi\* - bcmfxk@gwumc.edu

\* Corresponding author

Published: 09 August 2006

Received: 18 July 2006

Retrovirology 2006, 3:50 doi:10.1186/1742-4690-3-50

Accepted: 09 August 2006

This article is available from: <http://www.retrovirology.com/content/3/1/50>

© 2006 Pumfery et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

During interphase, each cell contains a single centrosome that acts as a microtubule organizing center for cellular functions in interphase and in mitosis. Centrosome amplification during the S phase of the cell cycle is a tightly regulated process to ensure that each daughter cell receives the proper complement of the genome. The controls that ensure that centrosomes are duplicated exactly once in the cell cycle are not well understood. In solid tumors and hematological malignancies, centrosome abnormalities resulting in aneuploidy is observed in the majority of cancers. These phenotypes are also observed in cancers induced by viruses, including adult T cell lymphoma which is caused by the human T cell lymphotropic virus Type I (HTLV-I). Several reports have indicated that the HTLV-I transactivator, Tax, is directly responsible for the centrosomal abnormalities observed in ATL cells. A recent paper in *Nature Cell Biology* by Ching *et al.* has shed some new light into how Tax may be inducing centrosome abnormalities. The authors demonstrated that 30% of ATL cells contained more than two centrosomes and expression of Tax alone induced supernumerary centrosomes. A cellular coiled-coil protein, TaxIBP2, was shown to interact with Tax and disruption of this interaction led to failure of Tax to induce centrosome amplification. Additionally, down-regulation of TaxIBP2 led to centrosome amplification. These results suggest that TaxIBP2 may be an important block to centrosome re-duplication that is observed in normal cells. Presently, a specific cellular protein that prevents centrosome re-duplication has not been identified. This paper has provided further insight into how Tax induces centrosome abnormalities that lead to ATL. Lastly, additional work on TaxIBP2 will also provide insight into how the cell suppresses centrosome re-duplication during the cell cycle and the role that TaxIBP2 plays in this important cellular pathway.

### Background

Faithful duplication of the genetic content of cells and proper segregation into two daughter cells are two highly regulated and distinct processes. The cell cycle determines when cells will duplicate their genome and the four phases are regulated by a series of different cyclin/cyclin-dependent kinase complexes. Duplication of the genome

occurs during the S phase while segregation of duplicated chromosomes into daughter cells occurs during the M (mitotic) phase. Proper segregation of chromosomes during M phase requires that the centrosomes are faithfully duplicated prior to mitosis. Duplication of centrosomes occurs at the end of G1 and the beginning of the S phase, yet is regulated separately from the duplication of the

genome. Dysregulation of either the cell cycle or centrosome duplication can result in major genetic alterations and cancer [1-3]. Changes in centrosome duplication have been observed in cancers caused by genetic mutations as well as in cancers induced by viruses. A recent report by Ching *et al.* [4] further elucidates how the HTLV-1 Tax protein may modulate centrosome duplication resulting in aneuploidy and the development of adult T cell leukemia (ATL).

The centrosome functions as the microtubule organizing center (MTOC) of the cell and modulates the microtubule network critical for chromosome segregation, cell division, intracellular transport, and development [1-3]. The centrosome is composed of a pair of centrioles surrounded by the pericentriolar material [1,2], which is composed of at least 50 different proteins including several different tubulins, centrin, and proteins containing a coiled-coil motif [2]. To prepare for chromosome segregation in the next M phase, a single centrosome needs to be duplicated. During the G0 and G1 phases of the cell cycle there is a single centrosome that is inherited from the mother cell during the previous cell division, which is duplicated during the late G1 and S phases. During mitosis, the number of spindle poles is generally determined by the number of centrosomes [1].

Centrosome duplication occurs in a semi-conservative manner and consists of four phases: (i) centriole splitting in which the mother and daughter centrioles detach; (ii) semiconservative centrosome duplication, where a new centriole is formed adjacent to the original centriole; (iii) maturation involves recruitment of pericentriolar material proteins; and (iv) centrosome separation during mitosis to form the spindle poles [1,2]. Centrosome duplication is regulated by a series of phosphorylation and dephosphorylation events mediated by several kinases including Cdk2, Nek2, polo-like kinases, and aurora kinases [1,2,5]. The coordination between replication and centrosome duplication suggests that these events may be regulated by a common mechanism. Cdk2 is attractive as the central

modulator since it is involved in both processes. Cdk2 is activated by binding to Cyclin E or Cyclin A at the start of S phase to induce DNA replication. Additionally, Cdk2 is required for centrosome duplication in a variety of cell systems [2,6]. Centrosomes are duplicated only once during each cell cycle and there is an apparent intrinsic block to reduplication of centrosomes [7]. However, the nature of this block is not known, but it apparently is overcome in cancer. Increased centrosome numbers leads to disturbances in mitosis and cytokinesis leading to errors in chromosome segregation and chromosome instability [6].

Alterations in centrosome duplication and the resulting aneuploidy are observed in a significant number of cancers. There are several possible mechanisms of centrosome amplification, including: (i) centrosome duplication more than once in a cell cycle; (ii) failure to undergo cytokinesis resulting in doubling of the genome [6]; and improper splitting, or fragmentation, of centrosomes in the M phase due to DNA damage [8]. Centrosome amplification is frequently seen in a number of solid cancers including breast, lung, colon, prostate, testicular, and head and neck squamous cell carcinoma [6], as well as several hematological malignancies such as Hodgkin's lymphoma, multiple myeloma, and chronic lymphocytic leukemia [9]. Some of the common genetic alterations in cancer associated with centrosome amplification include inactivation of p53, over-expression of Cyclin E [6], and mutation of BRCA1 [6,10].

Centrosome abnormalities are also observed in cancers caused by tumor viruses such as hepatitis B virus (HBV) [11-13], human papillomavirus [14,15], Kaposi's sarcoma-associated herpesvirus (KSHV) [16,17], and human T cell lymphotropic virus type 1 (HTLV-1) [4,18-20] [Table 1]. The HBV X protein induces centrosome amplification by activating the Ras-MEK-MAP kinase pathway [13]. Additionally, HBV X targets the Crm1 nuclear receptor, sequestering it in the cytoplasm [12]. Inactivation of Crm1 leads to abnormal centrioles and the presence of

**Table 1: Effect of various viral proteins on centrosomal abnormalities**

Viral Protein	Cellular Interacting Protein	Function in Normal Cells	Phenotype
HBV X	Crm1	Nuclear export receptor	Abnormal centrioles >2 centrosomes per cell
HPV E7	Rb	Tumor suppressor Transcriptional repressor	>2 centrosomes per cell
HPV E6	P53	Tumor suppressor	>2 centrosomes per cell
KSHV LANA	P53	Tumor suppressor	Abnormal mitotic spindles >2 centrosomes per cell
HTLV Tax	TaxBP181 RanBP1 Tax1BP2	Mitotic spindle checkpoint protein Regulator of Ran-GTPase pathway Regulates centrosome duplication	Multinucleated cells >2 centrosomes per cell >2 centrosomes per cell

more than two centrosomes in 39% of mitotic cells [11]. For human papillomavirus, both the E7 and E6 proteins contribute to supernumerary centrosomes. E7 inactivates Rb [14] and induces an increase in the number of centrioles, a portion of which give rise to mature centrosomes [15], whereas E6 inactivates p53 [14]. Expression of either protein alone will induce centrosome amplification; however, co-expression of E6 and E7 results in a higher incidence of supernumerary centrosomes [3,14]. Similar to HPV E6, the KSHV latency associated nuclear antigen (LANA) binds to p53 and inhibits the ability of p53 to transactivate cellular genes resulting in abnormal centrosomes, multinuclear cells, and other genomic abnormalities [17].

### Discussion

The HTLV-1 Tax protein contributes to the ability of the virus to cause ATL by inactivating p53 [21-24], altering the cell cycle [25,26], and inactivating Rb [27]. As Tax is able to deregulate cellular checkpoints, most notably the G1/S transition [28,29], it is not surprising that this oncoprotein is involved in mitotic disruption. An early report demonstrated that expression of Tax alone was sufficient to induce both centric (containing a kinetochore and representative of segregation defects) and acentric (representative of DNA damage) micronuclei [30]. While clastogenic effects induced by Tax appears to be due to subversion of DNA damage repair pathways [31-34], Tax involvement in aneuploid damage was less clear. A study by Jin *et al.* [20] indicated that the association of Tax with hsMAD1, a mitotic spindle checkpoint (MSC) protein, led to the translocation of both MAD1 and MAD2 to the cytoplasm. The MSC is necessary to halt the cell cycle when an error(s) in chromosome segregation is detected. When all of the kinetochores are attached to both poles of the mitotic spindle apparatus and chromosomes are equally segregated to opposite poles, the block is then released [35]. By disrupting normal localization of these factors (*i.e.* MAD1 and MAD2) to kinetochores following chromosomal missegregation, Tax was able to usurp this checkpoint and thus allow for accumulation of multinucleated cells, a common phenotype of ATL cells. However, several questions remained including (i) was aneuploidogenic damage simply the accumulation of natural occurring mis-segregation events [36] due to Tax or (ii) were there other Tax-related mechanisms either direct or indirect, that induced aneuploid damage?

Evidence for the direct involvement of Tax in aneuploidy development was recently shown by Peloponese and colleagues [19]. Tax was shown to localize to centrosomes during mitosis and interact with the Ran-GTPase pathway through the Ran-binding protein 1 (RanBP1). This protein complex regulates mitotic centrosome stability and microtubule nucleation and spindle formation [37,38].

Interaction of Tax with RanBP1 was shown to be necessary for Tax targeting to centrosomes and induction of supernumerary centrosomes. The interaction of Tax with the Ran-GTPase pathway to dysregulate centrosome duplication is reminiscent of HBV X interaction with Crm1 [12], a Ran-GTP binding nuclear export receptor [39].

The involvement of Tax in centrosome amplification has been further strengthened by recent findings from Ching *et al.* [4]. Tax was observed to localize to centrosomes in both transfected cells and HTLV-1 transformed cells. About 30% of ATL cells displayed centrosome amplification (>2 centrosomes) and 30-80% of cells expressing Tax alone had supernumerary centrosomes indicating that Tax was responsible for the abnormal centrosome phenotype in ATL. Through a yeast two-hybrid approach, Tax was shown to associate with a coiled-coil protein, Tax1BP2 (formerly known as TaxBP121). Tax1BP2 has high sequence similarity to C-Nap1 (centrosomal Nek2-associated protein 1), a protein involved in centrosome cohesion during the interphase of cell cycle [40], and was shown to be part of the centrosome complex. Tax interaction with Tax1BP2 was shown to be necessary for Tax to induce supernumerary centrosomes and over-expression of Tax1BP2 countered this effect. Additionally, down-regulation of Tax1BP2 expression by siRNAs resulted in amplification of centrosomes leading the authors to speculate that Tax1BP2 may be the intrinsic block to re-duplication proposed by Wong and Stearn [7]. This interaction provides a novel way for Tax to induce aneuploidogenic damage through centrosome amplification. Furthermore, if Tax1BP2 proves to be important for the block to centrosome duplication this will be the first protein identified to perform such a function. However, several questions remain: (i) is there differential expression or localization of Tax1BP2 during the cell cycle; (ii) does knock-down of Tax1BP2 expression in the S and G2 phases allow for centrosome re-duplication; and more importantly (iii) how does Tax1BP2 function to limit amplification of centrosomes?

### Conclusion

Centrosome amplification, which results in aneuploidy, is seen in a large number of cancers, including adult T cell lymphoma, and the HTLV-1 transactivator Tax plays a critical role in this process. Tax targets several pathways to induce uncontrolled cell growth including Rb [27], cyclin D2 [25,26,41-43], and Cdk 4 [44-48]. However, development of ATL takes decades and only a small percentage of infected individuals' progress to ATL, indicating that additional genetic alterations are required. Interestingly, while lymphoma and acute stage ATL patients display a high amount of structural (clastogenic) and numerical (aneuploid) damage, earlier stage patients (smoldering/chronic) do exhibit some aneuploid damage as exhibited

by multinucleated T cells. This suggests that aneupenic damage is an early phenomenon in ATL development; however, how aneupenic damage contributes to malignant transformation is still not known. Duplication of centrosomes and proper segregation of chromosomes into daughter cells is a highly regulated process involving a number of centrosome proteins, which have yet to be fully elucidated. Recent work has demonstrated that Tax can induce genomic instability and aneuploidy by directly targeting several proteins involved in centrosome duplication [Figure 1]. Interaction with Tax1BP2 may interrupt the intrinsic block to centrosome reduplication allowing for centrosome amplification. As mentioned by the authors, future studies will entail determining the mechanism of Tax1BP2 contributing to this block and how Tax may modulate this effect. Since Tax1BP2 appears to be phosphorylated and possibly regulated by Cdk2, Tax may bring Cdk2 into contact with Tax1BP2. Interaction with members of the APC induces a delay in mitotic entry and increased chromosomal instability. The centrosome amplification and chromosomal instabilities would allow for additional genetic alterations. Lastly, the interaction of Tax with Mad1, a component of the mitotic spindle checkpoint, inactivates this critical checkpoint allowing cells with supernumerary chromosomes and improperly segregated chromosomes to move through mitosis and into the next cell cycle. Additional questions remain as to whether there are Tax-dependent defects within cytokinesis that contribute to aneupenic damage and how the three Tax binding proteins identified by this group function together to regulate centrosome duplication. The role of Tax as a transcriptional regulator may also provide avenues for Tax to disrupt mitotic processes. Future studies will not only allow for better understanding of Tax involvement at mitosis but help to further elucidate how deregulation of this phase of the cell cycle contributes to carcinogenesis.

### Competing interests

The author(s) declare that they have no competing interests.

### Authors' contributions


AP and CF did the background research and most of the writing. FK edited and contributed to the figure development.

### Acknowledgements

This work was supported by grants from the George Washington University REF funds to A. Vertes and F. Kashanchi, and NIH grants AI44357, AI43894 and I3969 to F.K.

### References

- Hinchcliffe EH, Sluder G: "It takes two to tango": Understanding how centrosome duplication is regulated throughout the cell cycle. *Genes Dev* 2001, **15**:1167-1181.



**Figure 1**

### Role of Tax binding proteins in centrosomal abnormalities observed in ATL cells. (A) TaxBP181 (huMAD1).

During interphase TaxBP181 (light blue) is localized to the nucleus (yellow) and some co-localize with Tax (green) in infected cells. During the transition to prophase (1) in normal cells, TaxBP181 localizes to the kinetochores of chromosomes and allows proper segregation. (2) In normal cells during anaphase and telophase, TaxBP181 localizes to the midbody and finally in newly formed progeny nuclei. (3) When Tax is expressed in ATL cells, TaxBP181 is translocated to the nucleus, allowing for chromosome missegregation, resulting in multinucleated cells (4). (B) RanBP1. During interphase, Tax (green) is found in the nucleus and a portion is co-localized with RanBP1 (purple) on centrosomes. Interaction of Tax with RanBP1 dysregulates the centrosome duplication pathway resulting in cells with two or more centrosomes following mitosis. (C) Tax1BP2. During interphase Tax (green) is expressed in the nucleus and a portion co-localizes with pericentrin (purple) in centrosomes. During S phase centrosomes are duplicated. However, the interaction of Tax with Tax1BP2 (yellow) disrupts the normal controls that prevent centrosome re-duplication resulting in cells with more than two centrosomes. During mitosis, supernumerary centrosomes are separated into two daughter cells resulting in a percentage of cells with two or more centrosomes.

2. Wang Q, Hirohashi Y, Furuuchi K, Zhao H, Liu Q, Zhang H, Murali R, Berezhov A, Du X, Li B, Greene MI: **The centrosome in normal and transformed cells.** *DNA Cell Biol* 2004, **23**:475-489.
3. Doxsey S, McCollum D, Theurkauf W: **Centrosomes in cellular regulation.** *Annu Rev Cell Dev Biol* 2005, **21**:411-434.
4. Ching Y, Chan S, Jeang K, Jin D: **The retroviral oncoprotein tax targets the coiled-coil centrosomal protein TAXIBP2 to induce centrosome overduplication.** *Nat Cell Biol* 2006, **8**:717-724.
5. Hinchcliffe EH, Sluder G: **Centrosome duplication: Three kinases come up a winner!** *Curr Biol* 2001, **11**:R698-701.
6. Fukasawa K: **Centrosome amplification, chromosome instability and cancer development.** *Cancer Lett* 2005, **230**:6-19.
7. Wong C, Stearns T: **Centrosome number is controlled by a centrosome-intrinsic block to reduplication.** *Nat Cell Biol* 2003, **5**:539-544.
8. Hut HMJ, Lemstra W, Blaauw EH, Van Cappellen, Gert WA, Kampinga HH, Sibon OCM: **Centrosomes split in the presence of impaired DNA integrity during mitosis.** *Mol Biol Cell* 2003, **14**:1993-2004.
9. Krämer A, Neben K, Ho AD: **Centrosome aberrations in hematological malignancies.** *Cell Biol Int* 2005, **29**:375-383.
10. Deng C: **Roles of BRCA1 in centrosome duplication.** *Oncogene* 2002, **21**:6222-6227.
11. Forgues M, Difilippantonio MJ, Linke SP, Ried T, Nagashima K, Feden J, Valerie K, Fukasawa K, Wang XW: **Involvement of Crm1 in hepatitis B virus X protein-induced aberrant centriole replication and abnormal mitotic spindles.** *Mol Cell Biol* 2003, **23**:5282-5292.
12. Forgues M, Marrogi AJ, Spillare EA, Wu CG, Yang Q, Yoshida M, Wang XW: **Interaction of the hepatitis B virus X protein with the Crm1-dependent nuclear export pathway.** *J Biol Chem* 2001, **276**:22797-22803.
13. Yun C, Cho H, Kim S, Lee J, Park SY, Chan GK, Cho H: **Mitotic aberration coupled with centrosome amplification is induced by hepatitis B virus X oncoprotein via the ras-mitogen-activated protein/extracellular signal-regulated kinase-mitogen-activated protein pathway.** *Mol Cancer Res* 2004, **2**:159-169.
14. Duensing S, Münger K: **Centrosomes, genomic instability, and cervical carcinogenesis.** *Crit Rev Eukaryot Gene Expr* 2003, **13**:9-23.
15. Duensing S, Münger K: **The human papillomavirus type 16 E6 and E7 oncoproteins independently induce numerical and structural chromosome instability.** *Cancer Res* 2002, **62**:7075-7082.
16. Pan H, Zhou F, Gao S: **Kaposi's sarcoma-associated herpesvirus induction of chromosome instability in primary human endothelial cells.** *Cancer Res* 2004, **64**:4064-4068.
17. Si H, Robertson ES: **Kaposi's sarcoma-associated herpesvirus-encoded latency-associated nuclear antigen induces chromosomal instability through inhibition of p53 function.** *J Virol* 2006, **80**:697-709.
18. Nitta T, Kanai M, Sugihara E, Tanaka M, Sun B, Sonoda S, Miwa M, Nagasawa T: **Centrosome amplification in adult T-cell leukemia and human T-cell leukemia virus type I tax-induced human T cells.** *Cancer Sci* 2006 in press.
19. Peloponese J Jr, Haller K, Miyazato A, Jeang K: **Abnormal centrosome amplification in cells through the targeting of ran-binding protein-1 by the human T cell leukemia virus type-I tax oncoprotein.** *Proc Natl Acad Sci USA* 2005, **102**:18974-18979.
20. Jin DY, Spencer F, Jeang KT: **Human T cell leukemia virus type I oncoprotein tax targets the human mitotic checkpoint protein MAD1.** *Cell* 1998, **93**:81-91.
21. Tabakin-Fix Y, Azran I, Schavinky-Khrapunsky Y, Levy O, Aboud M: **Functional inactivation of p53 by human T-cell leukemia virus type I tax protein: Mechanisms and clinical implications.** *Carcinogenesis* 2006, **27**:673-681.
22. Grassmann R, Aboud M, Jeang K: **Molecular mechanisms of cellular transformation by HTLV-I tax.** *Oncogene* 2005, **24**:5976-5985.
23. Jeong S, Pise-Masison CA, Radonovich MF, Park HU, Brady JN: **Activated AKT regulates NF-kappaB activation, p53 inhibition and cell survival in HTLV-I-transformed cells.** *Oncogene* 2005, **24**:6719-6728.
24. Pise-Masison CA, Brady JN: **Setting the stage for transformation: HTLV-I tax inhibition of p53 function.** *Front Biosci* 2005, **10**:919-930.
25. Kehn K, Berro R, de la Fuente C, Strouss K, Ghedin E, Dadgar S, Bottazzi ME, Pumfery A, Kashanchi F: **Mechanisms of HTLV-I transformation.** *Front Biosci* 2004, **9**:2347-2372.
26. Kehn K, Deng L, de la Fuente C, Strouss K, Wu K, Maddukuri A, Baylor S, Rufner R, Pumfery A, Bottazzi ME, Kashanchi F: **The role of cyclin D2 and p21/waf1 in human T-cell leukemia virus type I infected cells.** *Retrovirology* 2004, **1**:6-6.
27. Kehn K, Fuente Cdl, Strouss K, Berro R, Jiang H, Brady J, Mahieux R, Pumfery A, Bottazzi ME, Kashanchi F: **The HTLV-I tax oncoprotein targets the retinoblastoma protein for proteasomal degradation.** *Oncogene* 2005, **24**:525-540.
28. Mulloy JC, Kislyakova T, Cereseto A, Casareto L, LoMonico A, Fullen J, Lorenzi MV, Cara A, Nicot C, Giam C, Franchini G: **Human T-cell lymphotropic/leukemia virus type I tax abrogates p53-induced cell cycle arrest and apoptosis through its CREB/ATF functional domain.** *J Virol* 1998, **72**:8852-8860.
29. Schmitt I, Rosin O, Rohwer P, Gossen M, Grassmann R: **Stimulation of cyclin-dependent kinase activity and G1- to S-phase transition in human lymphocytes by the human T-cell leukemia/lymphotropic virus type I tax protein.** *J Virol* 1998, **72**:633-640.
30. Majone F, Semmes OJ, Jeang KT: **Induction of micronuclei by HTLV-I tax: A cellular assay for function.** *Virology* 1993, **193**:456-459.
31. Majone F, Jeang KT: **Clastogenic effect of the human T-cell leukemia virus type I tax oncoprotein correlates with unstabilized DNA breaks.** *J Biol Chem* 2000, **275**:32906-32910.
32. Marriott SJ, Lemoine FJ, Jeang K: **Damaged DNA and miscounted chromosomes: Human T cell leukemia virus type I tax oncoprotein and genetic lesions in transformed cells.** *J Biomed Sci* 2002, **9**:292-298.
33. Jeang K, Giam C, Majone F, Aboud M: **Life, death, and tax: Role of HTLV-I oncoprotein in genetic instability and cellular transformation.** *J Biol Chem* 2004, **279**:31991-31994.
34. Majone F, Luisetto R, Zamboni D, Iwanaga Y, Jeang K: **Ku protein as a potential human T-cell leukemia virus type I (HTLV-I) tax target in clastogenic chromosomal instability of mammalian cells.** *Retrovirology* 2005, **2**:45-45.
35. Chan GK, Liu S, Yen TJ: **Kinetochore structure and function.** *Trends Cell Biol* 2005, **15**:589-598.
36. Norppa H, Falck GC: **What do human micronuclei contain?** *Mutagenesis* 2003, **18**:221-233.
37. Zhang C, Hughes M, Clarke PR: **Ran-GTP stabilises microtubule asters and inhibits nuclear assembly in xenopus egg extracts.** *J Cell Sci* 1999, **112(Pt 14)**:2453-2461.
38. Di Fiore B, Ciciarello M, Mangiacasale R, Palena A, Tassin A, Cundari E, Lavia P: **Mammalian RanBP1 regulates centrosome cohesion during mitosis.** *J Cell Sci* 2003, **116**:3399-3411.
39. Arnaoutov A, Azuma Y, Ribbeck K, Joseph J, Boyarchuk Y, Karpova T, McNally J, Dasso M: **Crm1 is a mitotic effector of ran-GTP in somatic cells.** *Nat Cell Biol* 2005, **7**:626-632.
40. Mayor T, Stierhof YD, Tanaka K, Fry AM, Nigg EA: **The centrosomal protein C-Nap1 is required for cell cycle-regulated centrosome cohesion.** *J Cell Biol* 2000, **151**:837-846.
41. Mori N, Fujii M, Hinz M, Nakayama K, Yamada Y, Ikeda S, Yamasaki Y, Kashanchi F, Tanaka Y, Tomonaga M, Yamamoto N: **Activation of cyclin D1 and D2 promoters by human T-cell leukemia virus type I tax protein is associated with IL-2-independent growth of T cells.** *Int J Cancer* 2002, **99**:378-385.
42. Santiago F, Clark E, Chong S, Molina C, Mozafari F, Mahieux R, Fujii M, Azimi N, Kashanchi F: **Transcriptional up-regulation of the cyclin D2 gene and acquisition of new cyclin-dependent kinase partners in human T-cell leukemia virus type I-infected cells.** *J Virol* 1999, **73**:9917-9927.
43. Akagi T, Ono H, Shimotohno K: **Expression of cell-cycle regulatory genes in HTLV-I infected T-cell lines: Possible involvement of Tax1 in the altered expression of cyclin D2, p18Ink4 and p21Waf1/Cip1/Sdi1.** *Oncogene* 1996, **12**:1645-1652.
44. Li J, Li H, Tsai M: **Direct binding of the N-terminus of HTLV-I tax oncoprotein to cyclin-dependent kinase 4 is a dominant path to stimulate the kinase activity.** *Biochemistry* 2003, **42**:6921-6928.
45. Fraedrich K, Müller B, Grassmann R: **The HTLV-I tax protein binding domain of cyclin-dependent kinase 4 (CDK4)**

- includes the regulatory **PSTAIR** helix. *Retrovirology* 2005, **2**:54-54.
46. Li J, Li H, Tsai MD: **Direct binding of the N-terminus of HTLV-I tax oncoprotein to cyclin-dependent kinase 4 is a dominant path to stimulate the kinase activity.** *Biochemistry (N Y)* 2003, **42**:6921-6928.
  47. Haller K, Wu Y, Derow E, Schmitt I, Jeang KT, Grassmann R: **Physical interaction of human T-cell leukemia virus type I tax with cyclin-dependent kinase 4 stimulates the phosphorylation of retinoblastoma protein.** *Mol Cell Biol* 2002, **22**:3327-3338.
  48. Neuveut C, Low KG, Maldarelli F, Schmitt I, Majone F, Grassmann R, Jeang KT: **Human T-cell leukemia virus type I tax and cell cycle progression: Role of cyclin D-cdk and p110Rb.** *Mol Cell Biol* 1998, **18**:3620-3632.

Publish with **BioMed Central** and every scientist can read your work free of charge

*"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."*

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

