

Poster presentation

Open Access

HIV-1 natural viral suppressors: control of viral replication in the absence of therapy

Mohammad Sajadi*

Address: University of Maryland School of Medicine, Baltimore, Maryland, USA

* Corresponding author

from 2006 International Meeting of The Institute of Human Virology
Baltimore, USA. 17–21 November, 2006

Published: 21 December 2006

Retrovirology 2006, **3**(Suppl 1):P55 doi:10.1186/1742-4690-3-S1-P55

© 2006 Sajadi; licensee BioMed Central Ltd.

Background

HIV Natural Viral Suppressors (NVS) are individuals with HIV-1 that are able to suppress HIV viral replication to undetectable levels in the absence of antiretroviral therapy. By contrast, Long-Term Nonprogressors (LTNP), represent individuals who are HIV-1 infected but who maintain high CD4 counts over many years (no where in the definition of LTNPs is viral load considered, and in most series the HIV-1 viral loads range from 10^2 to 10^4 copies/ml) [1-3]. We present here a cohort of NVS, providing a case definition for identifying such individuals and the results of preliminary evaluations.

Methods

Inclusion/Exclusion Criteria:

1. Confirmed HIV-1 by Western blot.
2. Positive HIV-1 Proviral DNA.
3. Viral loads <400 copies/ml at all time points for at least a 2 year period and 4 viral load measurements (One viral

blip or viral load > 400 copies/ml during the above time period was permitted if the subsequent viral loads were <400).

4. No more than 2 weeks of antiretroviral (ART) use (unless administered for pregnancy prophylaxis)

Studies included High Input PCR for Proviral DNA copy number.

Results

35 patients were identified as NVS, and 22 have been enrolled. Demographics are shown in Table 1.

Using a high-input assay developed specifically for this cohort (limit of detection 1 copy/ 10^6 PBMCs), proviral DNA copy number ranged from 1–74 copies/ 10^6 PBMCs (refer to Figure 1). The mean proviral DNA copy number was lower than both the controls we used in this study, HIV progressors and individuals with low viral loads (Figure 1A). The later group consisted of HIV infected individuals who had viral loads of 500–15,000 copies/ml, viral

Table 1: Demographics of the NVS cohort

Age	51 (range 38–60)
Sex	Male 64%, Female 36%
Race	African American 100%
Median years with HIV	12 (range 2–19)
Median years viral suppression	5 (range 2–9)
Median of latest CD4 count (cells/ul)	887 (range 346–1564)
Average number of viral loads tested	13 (range 4–21)

6. Vesanen M, Stevens C, Taylor P, et al.: **Stability in controlling viral replication identifies long-term nonprogressors as a distinct subgroup among HIV type 1-infected persons.** *J Virol* 1996, **70(12)**:9035-9040.
7. Brostrom C, Visco-Comandini U, Yun Z, Sonnerborg A: **A Longitudinal quantification of human immunodeficiency virus type 1 DNA and RNA in long-term nonprogressors.** *J Infect Dis* 1999, **179(6)**:1542-1548.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

