

Oral presentation

Open Access

## Potent cellular and humoral immunity against HIV-1 elicited in mice by a DNA-prime/MVA-boost vaccine regimen intended for human use

Britta Wahren\*<sup>1</sup>, Andreas Bråve<sup>1</sup>, Andreas Boberg<sup>1</sup>, Erik Rollman<sup>1</sup>, Deborah Birx<sup>2</sup>, Josephine Cox<sup>2</sup>, Merlin Robb<sup>2</sup>, Bernhard Moss<sup>2</sup>, Pontus Blomberg<sup>1</sup>, Gunnel Biberfeldt<sup>1</sup> and Eric Sandström<sup>1</sup>

Address: <sup>1</sup>Swedish Institute for Infectious Disease Control, Karolinska University Hospital and Karolinska Institute, Stockholm and <sup>2</sup>Walter Reed Army Institute of Research and National Institutes of Health, Silver Spring, Maryland, USA

Email: Britta Wahren\* - Britta.Wahren@smi.ki.se

\* Corresponding author

from 2006 International Meeting of The Institute of Human Virology  
Baltimore, USA. 17–21 November, 2006

Published: 21 December 2006

*Retrovirology* 2006, **3**(Suppl 1):S83 doi:10.1186/1742-4690-3-S1-S83

© 2006 Wahren et al; licensee BioMed Central Ltd.

In an experimental vaccine model, mice were primed three times with plasmids encoding multiple subtypes (A, B and C) of gag, envelope (env) and reverse transcriptase (RT) and adjuvant rGM-CSF followed by modified vaccinia Ankara (MVA) with the recombinant form A<sub>E</sub>, in theory providing protection against HIV subtypes A-E. The cellular responses, as measured by IFN-gamma secretion gave up to 2000 gag-specific SFC/million PBMC. The humoral and cellular responses were further increased by the MVA-boosts with env and gag specific ELISpot responses above 3500 secreting/10<sup>6</sup> cells. Intracellular cytokine staining showed remarkably high numbers (~15%) of gag and env specific CD8<sup>+</sup> T cells. This paved the way for our clinical trial with the multigene/multisubtype DNA plasmids boosted with the MVA construct. Conclusions: This preclinical study clearly shows the potential of combining these particular DNA and MVAs in a prime/boost regimen and that it is possible to induce a strong and broad humoral and cellular response directed against several parts of HIV as well as to several subtypes of the virus.