

Oral presentation

Open Access

TRIM5alpha contributes to the anti-viral state

Thomas Pertel^{1,2}, Martha R Neagu^{1,2,3}, Stephane Hausman²,
Madeleine Zufferey², Pradeep Uchil⁴, Walther Mothes⁴ and
Jeremy Luban^{*1,2,3}

Address: ¹Institute for Research in Biomedicine, Bellinzona, Switzerland, ²Department of Microbiology & Molecular Medicine, University of Geneva, Geneva, Switzerland, ³Department of Microbiology, Columbia University, New York, USA and ⁴Microbial Pathogenesis, Yale University, New Haven, USA

* Corresponding author

from *Frontiers of Retrovirology: Complex retroviruses, retroelements and their hosts*
Montpellier, France. 21-23 September 2009

Published: 24 September 2009

Retrovirology 2009, **6**(Suppl 2):O6 doi:10.1186/1742-4690-6-S2-O6

This abstract is available from: <http://www.retrovirology.com/content/6/S2/O6>

© 2009 Pertel et al; licensee BioMed Central Ltd.

TRIM5alpha is a host restriction factor that mediates a CA-specific block to retroviral infection. TRIM5alpha binding to the viral protein is only detectable when CA is in the multimeric protein lattice of the retroviral core. This suggested that TRIM5alpha might function as a pattern recognition receptor (PRR) analogous to a TLR. Additionally, *TRIM5* is located within a cluster of interferon-stimulated genes. We therefore sought evidence for functional links between TRIM5alpha and innate immune factors.

TRIM5alpha was induced by type I interferons and PRR agonists in THP-1 cells and monocyte-derived dendritic cells and macrophages. Induction kinetics, knockdown of IRF3, STAT1, STAT2, or IFN α R, and IFN α R-blocking antibodies, demonstrated that TRIM5alpha induction is ISGF3-dependent. In parallel, we found that establishment of an antiviral state with interferon or PRR agonists prevented HIV-1 transduction of THP1 cells or monocyte-derived dendritic cells and macrophages. TRIM5alpha knockdown in THP1 cells largely rescued HIV-1 from the antiviral state, independent of envelope pseudotype. This effect of TRIM5 knockdown was independent of CA and even observed with non-retroviruses. We have now established that TRIM5 contributes to the antiviral state via effects on specific factors that act upstream of NF κ B and AP-1.