Retrovirology



Poster presentation

Open Access

Inactivation of cell associated-HIV-I in breast milk by treatment with the alkyl sulfate microbicide sodium dodecyl sulfate (SDS)

Edouard Tuaillon*^{1,2}, Kuda Mutasa³, Pierre-Alain Rubbo¹, Laura Choteau², Florence Naudan², Karine Bollore¹, Jean-Pierre Vendrell¹,² and Philippe Van de Perre¹,²

Address: ¹Université Montpellier 1, EA 4205 Transmission, Pathogenèse et Prévention de l'Infection par le VIH, France, ²Laboratoire de Bactériologie-Virologie, 191 Avenue Doyen Giraud, 34295 Montpellier, France and ³ZVITAMBO project, 1 Borrowdale Road, Harare, Zimbabwe * Corresponding author

from Frontiers of Retrovirology: Complex retroviruses, retroelements and their hosts Montpellier, France. 21-23 September 2009

Published: 24 September 2009

Retrovirology 2009, 6(Suppl 2):P85 doi:10.1186/1742-4690-6-S2-P85

This abstract is available from: http://www.retrovirology.com/content/6/S2/P85 © 2009 Tuaillon et al; licensee BioMed Central Ltd.

Background

Breast milk is recognized as a predominant mode of HIV-1 infection in infants. Cell-associated HIV-1 may be the main source of virus transmission during early phases of breast-feeding. We have previously observed that HIV-1-infected cells spontaneously producing virus persist in breast-milk from women under antiretroviral therapy. Treatment of expressed milk with a microbicide such as Sodium dodecyl sulfate (SDS) is proposed as a simple and safe option to inactivate both cell free and cell associated HIV-1 when formula feeding is not practicable. However, the effect of SDS on spontaneously HIV-1-producing CD4+T cells in breast milk has not been fully explored.

Materials and methods

In this report human milk was spiked by HIV-1-infected cells and treated with increasing exposure time and SDS concentration. CD4+T cell apoptosis and death, cell-associated HIV-1 RNA production, and spontaneous HIV-1-Ag cell secretion were quantified after SDS treatment.

Results

Cell death increases in presence of SDS in a concentrationand time-dependent manner, 50% of T lymphocytes death after 2 minutes with 0.14% SDS and 90% after 10 minutes with 0.1% SDS (Fig. 1A). Undetectable HIV-1 RNA cell production was achieved following exposure with a minimum concentration of 0.1% SDS during 2 minutes, IC50 = 0.03% (Fig. 1B). The inhibition of HIV-1 Ag secretion was explored at a single cell level by ELISpot assay. Using this method inactivation was 100% for SDS concentrations \geq 0.25% within 2 min (Fig. 1C).

Conclusion

By comparison with results previously reported using an infectivity model based on β -galactosidase MAGI cells¹, we observed that a two fold higher SDS concentration was required to complete inactivation of HIV-1-Ag-secreting cells. This concentration remains in the reported safe limits for ingestion of SDS by children (1 g/kg/day). Regarding the possible occurrence of transmission to the infant after controlling cell-free virus in breast milk from women on antiretroviral therapy, SDS treatment of expressed breast milk may be an interesting strategy to optimized the prevention of HIV-1 pediatric transmission.




Figure I

Effect of SDS exposure on T cell death (A), cell-associated HIV-I RNA production (B), and sponataeous HIV-I-Ag cell secretion (C).

References

 Urdaneta S, Wigdahl B, Neely EB, Berlin CM Jr, Schengrund CL, Lin HM, Howett MK: Inactivation of HIV-1 in breast milk by treatment with the alkyl sulfate microbicide sodium dodecyl sulfate (SDS). Retrovirology 2005, 2:28.

Publish with **Bio Med Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- \bullet yours you keep the copyright

Submit your manuscript here: http://www.biomedcentral.com/info/publishing_adv.asp

