

Poster presentation

Open Access

## P04-02. Increased breadth and potency of the neutralizing antibody response among dually-HIV-1-infected individuals

RL Powell\*<sup>1</sup>, T Kinge<sup>2</sup> and P Nyambi<sup>1</sup>

Address: <sup>1</sup>Microbiology, New York University School of Medicine, New York, NY, USA and <sup>2</sup>Ministry of Public Health, Yaounde, Cameroon

\* Corresponding author

from AIDS Vaccine 2009  
Paris, France. 19–22 October 2009

Published: 22 October 2009

Retrovirology 2009, 6(Suppl 3):P30 doi:10.1186/1742-4690-6-S3-P30

This abstract is available from: <http://www.retrovirology.com/content/6/S3/P30>

© 2009 Powell et al; licensee BioMed Central Ltd.

### Background

Neutralizing antibodies (nAbs) generated during HIV-1 infection are insufficient to protect against a 2nd (dual) infection; yet, little is known regarding the effect of a 2nd infection on the anti-HIV-1 nAb response. Therefore, the objective of this study was to analyze the effect of dual infection by subtype-discordant HIV-1 strains on the potency and breadth of this nAb response.

### Methods

Two sequential plasma samples from 4 dually-intersubtype-infected subjects, obtained ~6 months before and >12 months after the 2nd infection was identified, were tested against 7 heterologous viruses (5 primary isolates and 2 Tier1 viruses) representing subtypes A1, B, F2, G, and CRF02\_AG in the GHOST cell neutralization assay. Additionally, 23 singly-infected control subjects matched for disease stage, CD4 counts, and time between samples were studied. Plasma was assayed at 1:80 dilution to compare each plasma pair; subsequently, plasma from dually-infected subjects and 6 control subjects were assessed for magnitude and specificity of neutralization using plasma serial dilutions, 1:20–1:640.

### Results

At 1:80 plasma dilution, 3 of the 4 dually-infected subjects' plasma obtained after dual infection exhibited significantly increased neutralization compared to initial plasma. Increases in percent neutralization for all 3 plasma pairs against all 7 heterologous viruses were significant ( $0.05 > p > 0.001$ ), with the 2nd sample exhibiting 65%–96% neutralization. No control plasmas exhibited

>30% neutralization of any of the heterologous viruses; thus, there was an overall significant difference between case and control subject-sets ( $p = 0.009$ ). Fifty percent neutralization titers (IC<sub>50</sub>) of plasma obtained after dual infection from the 3 dually-infected subjects exhibited ~6X–100X increases compared to the initial samples ( $0.05 > p > 0.0001$ ). No differences in the magnitudes of neutralization were observed for any of the singly-infected, control sample pairs.

### Conclusion

These data propose that re-infection by a subtype-discordant virus broadens the anti-HIV-1 immune response, suggesting that vaccines incorporating and/or boosting with subtype-discordant immunogens may generate significantly improved anti-HIV-1 immunity.