



RESEARCH

Open Access

# Trafficking of some old world primate TRIM5 $\alpha$ proteins through the nucleus

Felipe Diaz-Griffero<sup>1</sup>, Daniel E Gallo<sup>4</sup>, Thomas J Hope<sup>4</sup> and Joseph Sodroski<sup>2,3\*</sup>

## Abstract

**Background:** TRIM5 $\alpha$  and TRIMCyp are cytoplasmic proteins that bind incoming retroviral capsids and mediate early blocks to viral infection. TRIM5 proteins form cytoplasmic bodies, which are highly dynamic structures. So far, TRIM5 proteins have been found only in the cytoplasm of cells. Interestingly, other proteins from the TRIM family localize to the nucleus. Therefore, we tested the possibility that TRIM5 proteins traffic to the nucleus and the impact of this trafficking on retroviral restriction.

**Results:** Here we report that the TRIM5 $\alpha$  proteins of two Old World primates, humans and rhesus monkeys, are transported into the nucleus and are shuttled back to the cytoplasm by a leptomycin B-sensitive mechanism. In leptomycin B-treated cells, these TRIM5 $\alpha$  proteins formed nuclear bodies that also contained TRIM19 (PML). Deletion of the amino terminus, including the linker 1 (L1) region, resulted in TRIM5 $\alpha$  proteins that accumulated in nuclear bodies. Leptomycin B treatment of TRIM5 $\alpha$ -expressing target cells only minimally affected the restriction of retrovirus infection.

**Conclusions:** We discovered the ability of human and rhesus TRIM5 $\alpha$  to shuttle into and out of the nucleus. This novel trafficking ability of TRIM5 $\alpha$  proteins could be important for an as-yet-unknown function of TRIM5 $\alpha$ .

**Keywords:** Restriction factor intracellular localization, retrovirus, leptomycin B

## Background

Proteins of the tripartite motif (TRIM) family contain RING, B-Box and coiled-coil domains, and thus have been referred to as RBCC proteins [1]. Members of this family have been implicated in diverse processes such as cell proliferation, differentiation, development, oncogenesis and apoptosis [1,2]. TRIM proteins often self-associate and, when overexpressed, aggregate to form nuclear or cytoplasmic bodies [1].

TRIM5 $\alpha$  is a cytoplasmic protein that is capable of restricting retrovirus infection in a species-dependent manner [3]. Variation among TRIM5 $\alpha$  proteins in different primates accounts for the early, post-entry blocks to infection by particular retroviruses [3-7]. For example, TRIM5 $\alpha$  proteins of Old World monkeys block human immunodeficiency virus (HIV-1) infection [3-5,7], whereas TRIM5 $\alpha$  proteins of New World

monkeys block infection by simian immunodeficiency virus (SIV<sub>mac</sub>) [8]. TRIM5 $\alpha$  from humans (TRIM5 $\alpha$ <sub>hu</sub>) is not as potent in restricting HIV-1 infection as Old World monkey TRIM5 $\alpha$ , but TRIM5 $\alpha$ <sub>hu</sub> potently restricts other retroviruses, e.g., N-tropic murine leukemia virus (N-MLV) and equine infectious anemia virus (EIAV) [3,4,6-8]. Owl monkeys, a New World monkey species, are unusual in not expressing a TRIM5 $\alpha$  protein, but instead express TRIMCyp, in which the RBCC domains of TRIM5 are fused to a cyclophilin A moiety [9,10].

Variation in splicing of the *TRIM5* primary transcript leads to the expression of TRIM5 isoforms, designated  $\alpha$ ,  $\gamma$  and  $\delta$  [1]. The TRIM5 $\alpha$  isoform contains, in addition to the RING, B-box 2 and coiled-coil domains, a carboxy-terminal B30.2 (SPRY) domain. The B30.2 (SPRY) domain is essential for the antiretroviral activity of TRIM5 $\alpha$  [3]. In some cases, the differences in the ability of TRIM5 $\alpha$  proteins from various primate species to restrict particular retroviruses are determined by sequences in the B30.2 (SPRY) domain [11-19]. The B30.2 (SPRY) domain in TRIM5 $\alpha$  and the cyclophilin A

\* Correspondence: [joseph\\_sodroski@dfci.harvard.edu](mailto:joseph_sodroski@dfci.harvard.edu)

<sup>2</sup>Department of Cancer Immunology and AIDS, Dana-Farber Cancer Institute, Department of Pathology, Division of AIDS, Harvard Medical School, Boston, MA 02115, USA

Full list of author information is available at the end of the article

domain in TRIMCyp allow these restriction factors to bind specifically to particular retroviral capsids [9,20-24]. Additional sequences in the B-box 2 domain contribute to higher-order self-association of TRIM5 $\alpha$ , which allows higher avidity for the retroviral capsid [25-27]. TRIM5 $\alpha$  proteins aggregate on the incoming retroviral capsid [28]; and, by as-yet-uncertain mechanisms, decrease the stability of the capsid [23,27,29,30].

Some TRIM proteins localize in the nucleus of cells. One example is TRIM19 (promyelocytic leukemia (PML) protein), which is a major component of nuclear domain 10 (ND10) bodies [31-33]. TRIM19 has been shown to interfere with the replication of several DNA and RNA viruses [34-41]. Both TRIM19 and TRIM5 $\alpha$  can inhibit herpes simplex virus replication [34,40,41], and both proteins are induced by type I interferons [18,42,43]. Thus, both cytoplasmic (e.g., TRIM5 $\alpha$ ) and nuclear (e.g., TRIM19) TRIM proteins may be involved in innate resistance to viral infection.

Here we study the intracellular localization of different TRIM5 $\alpha$  proteins and TRIMCyp after treatment of cells with leptomycin B. Leptomycin B is a specific inhibitor of the nuclear export factor CRM1 (exportin 1), which is critical for the export of proteins carrying a nuclear export sequence [44-49]. We document that TRIM5 $\alpha_{\text{hu}}$  and TRIM5 $\alpha_{\text{rh}}$  are actively shuttling between the cytoplasm and nucleus. By contrast, TRIM5 $\alpha$  proteins from the squirrel monkey (a New World monkey) and the cow did not accumulate in the nucleus upon leptomycin B treatment. TRIMCyp from owl monkeys also localized in the cytoplasm upon treatment with leptomycin B. We investigated the contribution of the nuclear export of TRIM5 $\alpha$  to the antiretroviral activity of the protein.

## Results

### Leptomycin B treatment results in nuclear accumulation of some TRIM5 $\alpha$ proteins

During the course of studying TRIM5 $\alpha$ , we tested the effect of leptomycin B (LMB), a specific inhibitor of nuclear export [44-49], on TRIM5 $\alpha$  localization. As dogs do not express a functional TRIM5 protein [14], we initially studied the localization of different TRIM5 $\alpha$  variants in canine cells. LMB treatment of Cf2Th canine cells stably expressing TRIM5 $\alpha_{\text{hu}}$  or TRIM5 $\alpha_{\text{rh}}$  resulted in the accumulation of these proteins in the nucleus (Figure 1). Both proteins were found in nuclear bodies after LMB treatment. By contrast, TRIMCyp and the TRIM5 $\alpha$  proteins from cows and several species of New World monkeys (squirrel monkeys, spider monkeys, marmosets and tamarins) remained localized in the cytoplasm after LMB treatment. These results suggest that TRIM5 $\alpha_{\text{hu}}$  and TRIM5 $\alpha_{\text{rh}}$  shuttle into the nucleus

and require active transport via the CRM1 protein to achieve cytoplasmic localization.

### Rapid accumulation of TRIM5 $\alpha_{\text{hu}}$ and TRIM5 $\alpha_{\text{rh}}$ in the nucleus after LMB treatment


To understand the kinetics of TRIM5 $\alpha_{\text{rh}}$  movement into the nucleus, we performed time-lapse fluorescent microscopy using a HeLa cell line stably expressing a TRIM5 $\alpha_{\text{rh}}$ -yellow fluorescent protein (YFP) fusion. These experiments revealed that treatment of cells with LMB resulted in a rapid accumulation of TRIM5 $\alpha_{\text{rh}}$ -GFP in the nucleus (Figure 2). Nuclear bodies containing TRIM5 $\alpha_{\text{rh}}$ -GFP were evident by 2 hours following the initiation of LMB treatment.

### Nuclear TRIM5 $\alpha_{\text{hu}}$ and TRIM5 $\alpha_{\text{rh}}$ proteins localize to ND10 bodies with TRIM19

To examine whether TRIM5 $\alpha_{\text{rh}}$  localizes to the same ND10 bodies as TRIM19 after LMB treatment, LMB-treated human cells stably expressing TRIM5 $\alpha_{\text{rh}}$  were stained with antibodies directed against TRIM19 and the hemagglutinin (HA) epitope tag on TRIM5 $\alpha_{\text{rh}}$ . The nuclear TRIM5 $\alpha_{\text{rh}}$  colocalized with TRIM19 (Figure 3A). Gold-labeled antibodies directed against the HA epitope tag on TRIM5 $\alpha_{\text{rh}}$  were used to investigate the structure of the nuclear bodies. The TRIM5 $\alpha$ -directed antibodies formed ring-like structures similar in appearance to those previously described for TRIM19 in ND10 bodies (Figure 3B) [31,33].

### Localization of a TRIM5 $\alpha_{\text{rh}}$ -pyruvate kinase fusion protein

The diameter of the nuclear pore is approximately 0.9 nm, which allows globular proteins less than 60 kD to diffuse freely through the channel [50-52]. TRIM5 $\alpha$  proteins (approximately 55 kD) are close to this diffusion limit. Moreover, TRIM5 $\alpha$  forms a stable dimer [20,21]; however, we do not know if the majority of TRIM5 $\alpha$  molecules that enter the nucleus are monomers or dimers. In addition, the molecular shape of TRIM5 $\alpha$  is unknown. These uncertainties raised the possibility that TRIM5 $\alpha$  is actively transported into the nucleus. To test this possibility, TRIM5 $\alpha_{\text{rh}}$  was fused to pyruvate kinase (PK), which is normally a cytoplasmic protein [53] and to the green fluorescent protein (GFP) to create the GFP-PK-TRIM5 $\alpha_{\text{rh}}$  chimeric protein. The GFP-PK-TRIM5 $\alpha_{\text{rh}}$  protein and a control GFP-PK protein were transiently expressed in HeLa cells (Figure 4). Localization of these proteins was examined in untreated and LMB-treated cells (Figure 4). After a two-hour treatment with 10 nM LMB, the GFP-PK-TRIM5 $\alpha_{\text{rh}}$  protein was detected in both the nucleus and the cytoplasm. By contrast, the GFP-PK protein was detected only in the cytoplasm of untreated and LMB-treated cells. These results



are consistent with the active transport of TRIM5 $\alpha_{rh}$  to the nucleus.


#### Identification of TRIM5 $\alpha_{rh}$ regions modulating localization

Proteins that localize to the nucleus and shuttle to the cytoplasm often contain nuclear localization and nuclear export signals, respectively [44-48]. TRIM5 $\alpha_{hu}$  and TRIM5 $\alpha_{rh}$  lack an obvious nuclear localization signal [54,55], nor do they contain sequences motifs predicted to function as nuclear export signals [56]. To gain some insight into the TRIM5 $\alpha_{rh}$  sequences that modulate nuclear localization and export, a series of TRIM5 $\alpha_{rh}$  mutants with deletions in N-terminal components were studied. The TRIM5 $\alpha_{rh}$   $\Delta$ 12 and TRIM5 $\alpha$   $\Delta$ 60 proteins behaved like wild-type TRIM5 $\alpha_{rh}$  with respect to localization in untreated cells (Figure 5A and Table 1). However, in the LMB-treated cells, TRIM5 $\alpha_{rh}$   $\Delta$ 12 and TRIM5 $\alpha$   $\Delta$ 60 exhibited a bright, more diffuse pattern with fewer nuclear bodies when compared with wild-type TRIM5 $\alpha_{rh}$  (Figure 5A and Table 1). These results

indicate that neither the immediate TRIM5 $\alpha_{rh}$  N-terminus nor the RING domain significantly influence nuclear localization and export. By contrast, the TRIM5 $\alpha_{rh}$   $\Delta$ 93 mutant localized to nuclear bodies and to the cytosol, even in the absence of LMB treatment (Figure 5B and Table 1). This localization pattern did not change significantly upon LMB treatment. Thus, deletion of TRIM5 $\alpha_{rh}$  sequences between residues 60 and 93, in the Linker 1 (L1) region of the protein, appears to decrease the efficiency of nuclear export of TRIM5 $\alpha_{rh}$ .

#### Contribution of nuclear export of TRIM5 $\alpha_{hu}$ and TRIM5 $\alpha_{rh}$ to retroviral restriction

To study the contribution of TRIM5 $\alpha$  nuclear export to retroviral restriction, we treated cells stably expressing TRIM5 $\alpha_{rh}$  and TRIM5 $\alpha_{hu}$  with LMB for two hours. Then the cells were challenged with recombinant HIV-1 and N-MLV expressing GFP. Treatment with LMB continued during the incubation of the cells with virus and




overnight thereafter. LMB treatment exerted only minimal effects on the ability of TRIM5 $\alpha_{rh}$  to restrict HIV-1 infection and on the ability of TRIM5 $\alpha_{hu}$  to inhibit N-MLV infection (Figure 6).

### Discussion

All characterized TRIM5 $\alpha$  proteins are located in the cytoplasm of expressing cells [15,28,57-59]. Here we report the surprising observation that some TRIM5 $\alpha$  proteins are imported into the nucleus and then exported back into the cytoplasm by a CRM1-dependent mechanism. Of interest, this transient routing through the nucleus was observed for the TRIM5 $\alpha$  proteins of two Old World primates, and not for the TRIM5 $\alpha$  proteins of a cow or several New World monkeys, or for the TRIMCyp protein of another New World monkey (the owl monkey). This raises the possibility that nuclear shuttling represents a property that was gained by Old World primate TRIM5 $\alpha$  proteins after the divergence from the New World monkeys.



Our results with the GFP-PK-TRIM5 $\alpha_{rh}$  fusion protein suggest that TRIM5 $\alpha_{rh}$  is actively transported into the nucleus, as the fusion protein is well above the size limit for passive diffusion of proteins through the nuclear pore [50-52]. Nonetheless, no typical nuclear localization motif is evident on TRIM5 $\alpha$  [54,55]. The accumulation of TRIM5 $\alpha_{hu}$  and TRIM5 $\alpha_{rh}$  in the nucleus after LMB treatment implicates a CRM1-dependent process in the export of these TRIM5 $\alpha$  proteins from the nucleus [44-49]. However, there are no classical nuclear export motifs in TRIM5 $\alpha$  proteins [56]. It is possible that TRIM5 $\alpha$  utilizes unusual motifs for interacting with nuclear pore proteins. Analysis of the localization of N-terminally truncated TRIM5 $\alpha_{rh}$  mutants suggests that deletion of residues 60-93, in the linker 1 (L1) region, disrupts the nuclear export of the protein. Whether this is a result of deletion of a non-canonical nuclear export signal or an indirect effect requires further investigation. As an example of the latter effect, the linker 1 (L1) regions could mediate the association




of TRIM5 $\alpha$ <sub>rh</sub> and TRIM5 $\alpha$ <sub>hu</sub> with another factor that shuttles between the nuclear and cytoplasm.

Despite the accumulation of TRIM5 $\alpha$ <sub>hu</sub> and TRIM5 $\alpha$ <sub>rh</sub> proteins in the nucleus after LMB treatment, restriction of N-MLV and HIV-1, respectively, remained

potent. Although it is possible that nuclear TRIM5 $\alpha$ <sub>hu</sub> and TRIM5 $\alpha$ <sub>rh</sub> can inhibit retrovirus infection, the specific recognition of the retroviral capsid, which does not enter the intact nucleus, is thought to be important for potent restriction [22,23]. A more likely explanation is



**Figure 4 Localization of a GFP-PK-TRIM5 $\alpha$  protein in leptomycin B-treated cells.** HeLa cells transiently expressing the fusion constructs GFP-PK or GFP-PK-TRIM5 $\alpha_{rh}$  were treated with 5 ng/ml of LMB or with the equivalent concentration of DMSO for 2 hours (A). Protein expression levels of the different fusion constructs were measured by Western blot using anti-GFP antibodies (B).



**Figure 5 Localization of TRIM5 $\alpha_{\text{rh}}$  N-terminal deletion mutants in leptomycin B-treated cells.** Cf2Th cells stably expressing wild-type TRIM5 $\alpha_{\text{rh}}$  or the indicated deletion mutant were treated with 5 ng/ml of LMB or DMSO for two hours. Treated cells were stained using anti-HA antibodies conjugated to FITC. TRIM5 $\alpha_{\text{rh}}$  domains are depicted for each variant, and the numbers of the amino acid residues at the boundaries of the different domains are shown (A). L1 represents the Linker 1 region. The TRIM5 $\alpha_{\text{rh}}\Delta 93$  protein bodies are located in the cellular nucleus (B). Cf2Th cells expressing TRIM5 $\alpha_{\text{rh}}\Delta 93$  were stained using anti-HA antibodies conjugated to FITC (green) and propidium iodide for nuclear staining (red). A representative image is shown.

**Table 1 Number of TRIM5 $\alpha$  cytoplasmic and nuclear bodies in LMB-treated cells**

	Number of cytoplasmic and nuclear bodies per 100 cells					
	DMSO			LMB		
	Cytoplasmic	Nuclear	Total	Cytoplasmic	Nuclear	Total
TRIM5 $\alpha_{rh}$	448	2	450	7	543	550
TRIM5 $\alpha_{rh}\Delta 12$	127	3	130*	5	128	133*
TRIM5 $\alpha_{rh}\Delta 60$	202	8	210*	2	78	80*
TRIM5 $\alpha_{rh}\Delta 93$	4	151	155	12	158	170

\*Cytoplasmic and nuclear bodies of TRIM5 $\alpha_{rh}\Delta 12$  and  $\Delta 60$  were on average larger than bodies observed for wt TRIM5 $\alpha_{rh}$  and TRIM5 $\alpha_{rh}\Delta 93$  proteins.

that the residual TRIM5 $\alpha$  protein in the cytoplasm of these overexpressing cells is sufficient to inhibit virus infection. Any newly synthesized TRIM5 $\alpha$  in these cells that has not yet entered the nucleus is potentially available for capsid interaction.

One caveat of these studies is the use of exogenously expressed TRIM5 $\alpha$  proteins to study nuclear shuttling. When better antibodies against endogenous TRIM5 $\alpha$  become available, the shuttling behavior of the endogenously expressed TRIM5 $\alpha$  protein can be examined.

What might be the possible advantage of having the Old World primate TRIM5 $\alpha$  proteins shuttle into and out of the nucleus? If these TRIM5 $\alpha$  proteins acquire post-translational modifications or binding partners in the process, our results suggest that such acquisition is apparently not necessary for HIV-1 or N-MLV restriction. The presence of TRIM5 $\alpha$  in the nucleus could be important for other TRIM5 $\alpha$  functions besides retroviral restriction. For example, Old World monkey TRIM5 $\alpha$  proteins have recently been shown to inhibit the infection of herpes simplex viruses 1 and 2 [41]. The colocalization of nuclear TRIM5 $\alpha$  in ND10 bodies with TRIM19, which also has anti-herpes virus activity [34,39,40], might have functional importance in this respect. Future studies should shed light on these interesting possibilities.

## Conclusions

Here we discovered the ability of human and rhesus TRIM5 $\alpha$  to shuttle into and out of the nucleus. Although not essential for retroviral restriction, this novel ability of TRIM5 $\alpha$  might be involved in other functions such as the ability of TRIM5 to trigger NF- $\kappa$ B [38].

## Methods

### Plasmid construction

The plasmids used to establish cell lines stably expressing TRIM5 $\alpha$  variants or TRIMCyp have been previously described [8,58]. The plasmids expressing mutant TRIM5 $\alpha_{rh}$  proteins with N-terminal deletions were constructed by polymerase chain reaction (PCR) amplification of *TRIM5* cDNA, as previously described

[3]. The amplified fragments were cloned into the EcoRI and Cla I sites of the pLPCX plasmid (Stratagene). All of the TRIM5 $\alpha$  proteins have an epitope tag from influenza hemagglutinin (HA). Human TRIM5 $\alpha$  has the HA tag at the carboxyl terminus, and all the other TRIM5 $\alpha$  proteins have the HA tag at the amino terminus.

### Creation of cells stably expressing TRIM5 $\alpha$ and TRIMCyp variants

Retroviral vectors encoding TRIM5 $\alpha$  or TRIMCyp proteins were created using the pLPCX vector plasmid [3]. Recombinant viruses were produced in 293T cells by cotransfecting the pLPCX plasmids with the pVPack-GP and pVPack-VSV-G packaging plasmids (Stratagene). The pVPack-VSV-G plasmid encodes the vesicular stomatitis virus (VSV) G envelope glycoprotein, which allows efficient entry into a wide range of vertebrate cells.


### Protein analysis

Cellular proteins were extracted with radioimmunoprecipitation assay (RIPA) buffer (10 mM Tris, pH 7.4; 100 mM NaCl; 1% sodium deoxycholate; 0.1% sodium dodecyl sulfate [SDS]; 1% NP-40; 2 mg of aprotinin/ml; 2 mg of leupeptin/ml; 1 mg of pepstatin A/ml; 100 mg of phenylmethylsulfonyl fluoride/ml). The cell lysates were analyzed by SDS-PAGE (10% acrylamide), followed by blotting onto nitrocellulose membranes (Amersham Pharmacia Biotech). Detection of protein by Western blotting utilized monoclonal antibodies that are specifically reactive with the HA epitope tag (Roche). Detection of proteins was performed by enhanced chemiluminescence (NEN Life Sciences Products).

### Infection with recombinant viruses expressing green fluorescent protein (GFP)

Recombinant HIV-1 or N-MLV expressing GFP were prepared as described [3]. HIV-1 viral stocks were quantified by measuring reverse transcriptase (RT) activity. For infections,  $3 \times 10^4$  HeLa human epithelial cells or Cf2Th canine cells seeded in 24-well plates were incubated in the presence of virus for 24 hours. Cells were washed and returned to culture for 48 hours, and then





**Figure 6 Effect of leptomycin B treatment of TRIM5 $\alpha$ -expressing cells on retrovirus restriction.** Cf2Th cells stably expressing TRIM5 $\alpha_{th}$  or transduced with the empty vector LPCX were challenged with increasing amounts of HIV-1-GFP in the presence of 5 ng/ml of LMB or DMSO (A). Similarly, Cf2Th cells stably expressing TRIM5 $\alpha_{hu}$  were challenged with increasing amounts of N-MLV-GFP in the presence of 5 ng/ml of LMB or DMSO (B). TE671 cells, which naturally express TRIM5 $\alpha_{hu}$ , were challenged with B-MLV-GFP and N-MLV-GFP in the presence of the indicated concentration of LMB or the DMSO control (C). The x-axis indicates the volume of a stock of recombinant GFP-expressing virus added to the target cells. Forty-eight hours after infection, the percentage of infected cells was measured by counting the GFP-positive cells using a flow cytometer. Similar results were obtained in three independent experiments.

subjected to FACS analysis with a FACScan (Becton Dickinson).

#### Intracellular location of TRIM5 variants

Localization of TRIM5 variants was studied as previously described [60]. Briefly, cells were grown overnight on 12-mm-diameter coverslips and fixed in 3.9% paraformaldehyde (Sigma) in phosphate-buffered saline (PBS; Cellgro) for 30 minutes. In some experiments, cells were incubated with 5 ng/ml leptomycin B (LMB) in medium for 2-10 hours prior to fixation. Cells were washed in PBS, incubated in 0.1 M glycine (Sigma) for 10 minutes, washed in PBS, and permeabilized with 0.05% saponin (Sigma) for 30 minutes. Samples were blocked with 10% donkey serum (Dako, Carpinteria, CA) for 30 minutes, and incubated for 1 hour with antibodies. HA-tagged proteins were stained using an anti-HA FITC-conjugated antibody, clone 3F10 (Roche). The TRIM19 (PML) protein was stained with an antibody against PML, sc-9863 (Santa Cruz Biotechnology, CA) and anti-goat Cy3-conjugated antibodies (Jackson ImmunoResearch, PA). Subsequently, samples were mounted for fluorescence microscopy by using the ProLong Antifade Kit (Molecular Probes, Eugene, OR). Images were obtained with a BioRad Radiance 2000 laser scanning confocal microscope with Nikon 60X N.A.1.4 optics.

#### Detection of TRIM5 $\alpha$ by electron microscopy

HeLa cells stably expressing HA-tagged TRIM5 $\alpha_{rh}$  treated with 5 ng/ml LMB for 2 h were removed from the tissue culture dish with 5 mM EDTA in PBS, pelleted, and resuspended in a small volume of 4% paraformaldehyde in 0.2 M sodium phosphate buffer, pH 7.4. Ultrathin sections were cut at -120°C with a cryo-diamond knife. Sections were picked up from the knife with a loop dipped in a 1:1 mixture of 2.3 M sucrose and 2% methylcellulose and transferred to a carbon-coated copper grid. Grids were left floating on PBS with the section facing down. Grids were washed in PBS and blocked in 1% bovine serum albumin (BSA) in PBS for 15 min. Grids were then incubated with the anti-HA 3F10 antibody (Roche) in 1% BSA in PBS for 30 min and washed four times for 15 min in PBS. Then, the grids were incubated with Protein A-gold 10-nm particles (Jackson ImmunoResearch) in 1% BSA in PBS for 20 min and washed four times for 15 min in PBS. Images were acquired using a transmission electron microscope JEOL 1200EX-80kV.

#### Acknowledgements

We thank Ms. Yvette McLaughlin and Ms. Elizabeth Carpelan for manuscript preparation and the National Institutes of Health (AI063987(JS), AI076094(JS), AI047770(TJH)) and a Center for AIDS Research Award AI60354), the International AIDS Vaccine Initiative, the Bristol-Myers Squibb Foundation,

and the late William F. McCarty-Cooper for research funding. F.D.-G. is a recipient of a K99/R00 Pathway to Independence Award from the National Institutes of Health (1K99MH086162-01), an American Foundation for AIDS Research Mathilde Krim fellowship in basic biomedical research (106987-43-RFHF), and a Claudia Adams Barr award from the Dana-Farber Cancer Institute. We also would like to thank the James B. Pendleton Charitable Trust. T.J.H. is funded by a P50 GM082545 from the NIH.

#### Author details

<sup>1</sup>Department of Microbiology and Immunology, Albert Einstein College of Medicine, Bronx, NY 10461, USA. <sup>2</sup>Department of Cancer Immunology and AIDS, Dana-Farber Cancer Institute, Department of Pathology, Division of AIDS, Harvard Medical School, Boston, MA 02115, USA. <sup>3</sup>Department of Immunology and Infectious Diseases, Harvard School of Public Health, Boston, MA 02115, USA. <sup>4</sup>Department of Cell and Molecular Biology, Northwestern University, Chicago, IL 60611, USA.

#### Authors' contributions

FDG designed and performed experiments, wrote the manuscript. DEG designed and performed experiments. TJH designed and performed experiments. JS designed experiments and wrote the manuscript. All authors read and approved the final manuscript.

#### Competing interests

The authors declare that they have no competing interests.

Received: 1 February 2011 Accepted: 15 May 2011

Published: 15 May 2011

#### References

1. Reymond A, Meroni G, Fantozzi A, Merla G, Cairo S, Luzi L, Riganelli D, Zanaria E, Messali S, Cainarca S, *et al*: The tripartite motif family identifies cell compartments. *Embo J* 2001, **20**:2140-2151.
2. Nisole S, Stoye JP, Saib A: TRIM family proteins: retroviral restriction and antiviral defence. *Nat Rev Microbiol* 2005, **3**:799-808.
3. Stremlau M, Owens CM, Perron MJ, Kiessling M, Autissier P, Sodroski J: The cytoplasmic body component TRIM5 $\alpha$  restricts HIV-1 infection in Old World monkeys. *Nature* 2004, **427**:848-853.
4. Hatzioannou T, Perez-Caballero D, Yang A, Cowan S, Bieniasz PD: Retrovirus resistance factors Ref1 and Lv1 are species-specific variants of TRIM5 $\alpha$ . *Proc Natl Acad Sci USA* 2004, **101**:10774-10779.
5. Keckesova Z, Ylilinen LM, Towers GJ: The human and African green monkey TRIM5 $\alpha$  genes encode Ref1 and Lv1 retroviral restriction factor activities. *Proc Natl Acad Sci USA* 2004, **101**:10780-10785.
6. Perron MJ, Stremlau M, Song B, Ulm W, Mulligan RC, Sodroski J: TRIM5 $\alpha$  mediates the postentry block to N-tropic murine leukemia viruses in human cells. *Proc Natl Acad Sci USA* 2004, **101**:11827-11832.
7. Yap MW, Nisole S, Lynch C, Stoye JP: Trim5 $\alpha$  protein restricts both HIV-1 and murine leukemia virus. *Proc Natl Acad Sci USA* 2004, **101**:10786-10791.
8. Song B, Javanbakht H, Perron M, Park DH, Stremlau M, Sodroski J: Retrovirus restriction by TRIM5 $\alpha$  variants from Old World and New World primates. *J Virol* 2005, **79**:3930-3937.
9. Nisole S, Lynch C, Stoye JP, Yap MW: A Trim5-cyclophilin A fusion protein found in owl monkey kidney cells can restrict HIV-1. *Proc Natl Acad Sci USA* 2004, **101**:13324-13328.
10. Sayah DM, Sokolskaja E, Berthoux L, Luban J: Cyclophilin A retrotransposition into TRIM5 explains owl monkey resistance to HIV-1. *Nature* 2004, **430**:569-573.
11. Song B, Gold B, O'Huigin C, Javanbakht H, Li X, Stremlau M, Winkler C, Dean M, Sodroski J: The B30.2(SPRY) domain of the retroviral restriction factor TRIM5 $\alpha$  exhibits lineage-specific length and sequence variation in primates. *J Virol* 2005, **79**:6111-6121.
12. Kono K, Bozek K, Domingues FS, Shioda T, Nakayama EE: Impact of a single amino acid in the variable region 2 of the Old World monkey TRIM5 $\alpha$  SPRY (B30.2) domain on anti-human immunodeficiency virus type 2 activity. *Virology* 2009, **388**:160-168.
13. Sebastian S, Grutter C, de Castillia CS, Pertel T, Olivari S, Grutter MG, Luban J: An invariant surface patch on the TRIM5 $\alpha$  PRYSPRY domain is required for retroviral restriction but dispensable for capsid binding. *J Virol* 2009, **83**:3365-3373.

14. Sawyer SL, Wu LI, Emerman M, Malik HS: Positive selection of primate TRIM5alpha identifies a critical species-specific retroviral restriction domain. *Proc Natl Acad Sci USA* 2005, **102**:2832-2837.
15. Perez-Caballero D, Hatzioannou T, Yang A, Cowan S, Bieniasz PD: Human tripartite motif 5alpha domains responsible for retrovirus restriction activity and specificity. *J Virol* 2005, **79**:8969-8978.
16. Ohkura S, Yap MW, Sheldon T, Stoye JP: All three variable regions of the TRIM5alpha B30.2 domain can contribute to the specificity of retrovirus restriction. *J Virol* 2006, **80**:8554-8565.
17. Yap MW, Nisole S, Stoye JP: A single amino acid change in the SPRY domain of human Trim5alpha leads to HIV-1 restriction. *Curr Biol* 2005, **15**:73-78.
18. Nakayama EE, Miyoshi H, Nagai Y, Shioda T: A specific region of 37 amino acid residues in the SPRY (B30.2) domain of African green monkey TRIM5alpha determines species-specific restriction of simian immunodeficiency virus SIVmac infection. *J Virol* 2005, **79**:8870-8877.
19. Stremmler M, Perron M, Welikala S, Sodroski J: Species-specific variation in the B30.2(SPRY) domain of TRIM5alpha determines the potency of human immunodeficiency virus restriction. *J Virol* 2005, **79**:3139-3145.
20. Kar AK, Diaz-Griffero F, Li Y, Li X, Sodroski J: Biochemical and biophysical characterization of a chimeric TRIM21-TRIM5alpha protein. *J Virol* 2008, **82**:11669-11681.
21. Langelier CR, Sandrin V, Eckert DM, Christensen DE, Chandrasekaran V, Alam SL, Aiken C, Olsen JC, Kar AK, Sodroski JG, Sundquist WI: Biochemical characterization of a recombinant TRIM5alpha protein that restricts human immunodeficiency virus type 1 replication. *J Virol* 2008, **82**:11682-11694.
22. Sebastian S, Luban J: TRIM5alpha selectively binds a restriction-sensitive retroviral capsid. *Retrovirology* 2005, **2**:40.
23. Stremmler M, Perron M, Lee M, Li Y, Song B, Javanbakht H, Diaz-Griffero F, Anderson DJ, Sundquist WI, Sodroski J: Specific recognition and accelerated uncoating of retroviral capsids by the TRIM5alpha restriction factor. *Proc Natl Acad Sci USA* 2006, **103**:5514-5519.
24. Diaz-Griffero F, Qin XR, Hayashi F, Kigawa T, Finzi A, Sarnak Z, Lienlaf M, Yokoyama S, Sodroski J: A B-box 2 Surface Patch Important for TRIM5 {alpha} Self-Association, Capsid-binding Avidity and Retrovirus Restriction. *J Virol* 2009.
25. Li X, Sodroski J: The TRIM5alpha B-box 2 domain promotes cooperative binding to the retroviral capsid by mediating higher-order self-association. *J Virol* 2008, **82**:11495-11502.
26. Diaz-Griffero F, Kar A, Lee M, Stremmler M, Poeschla E, Sodroski J: Comparative requirements for the restriction of retrovirus infection by TRIM5alpha and TRIMCyp. *Virology* 2007.
27. Campbell EM, Perez O, Anderson JL, Hope TJ: Visualization of a proteasome-independent intermediate during restriction of HIV-1 by rhesus TRIM5alpha. *J Cell Biol* 2008, **180**:549-561.
28. Diaz-Griffero F, Kar A, Perron M, Xiang SH, Javanbakht H, Li X, Sodroski J: Modulation of Retroviral Restriction and Proteasome Inhibitor-resistant Turnover by Changes in the TRIM5(alpha) B-box 2 Domain. *J Virol* 2007.
29. Diaz-Griffero F, Perron M, McGee-Estrada K, Hanna R, Maillard PV, Trono D, Sodroski J: A human TRIM5alpha B30.2/SPRY domain mutant gains the ability to restrict and prematurely uncoat B-tropic murine leukemia virus. *Virology* 2008.
30. Perron MJ, Stremmler M, Lee M, Javanbakht H, Song B, Sodroski J: The human TRIM5alpha restriction factor mediates accelerated uncoating of the N-tropic murine leukemia virus capsid. *J Virol* 2007, **81**:2138-2148.
31. Everett RD: DNA viruses and viral proteins that interact with PML nuclear bodies. *Oncogene* 2001, **20**:7266-7273.
32. Negorev D, Maul GG: Cellular proteins localized at and interacting within ND10/PML nuclear bodies/PODs suggest functions of a nuclear depot. *Oncogene* 2001, **20**:7234-7242.
33. Maul GG: Nuclear domain 10, the site of DNA virus transcription and replication. *Bioessays* 1998, **20**:660-667.
34. Everett RD, Maul GG: HSV-1 IE protein Vmw110 causes redistribution of PML. *Embo J* 1994, **13**:5062-5069.
35. Bjornal AS, Szekely L, Elgh F: Ebola virus infection inversely correlates with the overall expression levels of promyelocytic leukaemia (PML) protein in cultured cells. *BMC Microbiol* 2003, **3**:6.
36. Bonilla WV, Pinschewer DD, Klenerman P, Rousson V, Gaboli M, Pandolfi PP, Zinkernagel RM, Salvato MS, Hengartner H: Effects of promyelocytic leukemia protein on virus-host balance. *J Virol* 2002, **76**:3810-3818.
37. Djavani M, Rodas J, Lukashevich IS, Horejsh D, Pandolfi PP, Borden KL, Salvato MS: Role of the promyelocytic leukemia protein PML in the interferon sensitivity of lymphocytic choriomeningitis virus. *J Virol* 2001, **75**:6204-6208.
38. Tareen SU, Emerman M: Human Trim5alpha has additional activities that are uncoupled from retroviral capsid recognition. *Virology* 2011, **409**:113-120.
39. Kuroishi A, Bozek K, Shioda T, Nakayama EE: A single amino acid substitution of the human immunodeficiency virus type 1 capsid protein affects viral sensitivity to TRIM5 alpha. *Retrovirology* 2010, **7**:58.
40. Nakayama EE, Shioda T: Anti-retroviral activity of TRIM5 alpha. *Rev Med Virol* 2010, **20**:77-92.
41. Xu D, Holko M, Sadler AJ, Scott B, Higashiyama S, Berkofsky-Fessler W, McConnell MJ, Pandolfi PP, Licht JD, Williams BR: Promyelocytic leukemia zinc finger protein regulates interferon-mediated innate immunity. *Immunity* 2009, **30**:802-816.
42. Carthagen L, Parise MC, Ringard M, Chelbi-Alix MK, Hazan U, Nisole S: Implication of TRIM alpha and TRIMCyp in interferon-induced anti-retroviral restriction activities. *Retrovirology* 2008, **5**:59.
43. Nishi K, Yoshida M, Fujiwara D, Nishikawa M, Horinouchi S, Beppu T: Leptomycin B targets a regulatory cascade of crm1, a fission yeast nuclear protein, involved in control of higher order chromosome structure and gene expression. *J Biol Chem* 1994, **269**:6320-6324.
44. Watanabe M, Fukuda M, Yoshida M, Yanagida M, Nishida E: Involvement of CRM1, a nuclear export receptor, in mRNA export in mammalian cells and fission yeast. *Genes Cells* 1999, **4**:291-297.
45. Wolff B, Sanglier JJ, Wang Y: Leptomycin B is an inhibitor of nuclear export: inhibition of nucleocytoplasmic translocation of the human immunodeficiency virus type 1 (HIV-1) Rev protein and Rev-dependent mRNA. *Chem Biol* 1997, **4**:139-147.
46. Fornerod M, Ohno M, Yoshida M, Mattaj JW: CRM1 is an export receptor for leucine-rich nuclear export signals. *Cell* 1997, **90**:1051-1060.
47. Kuersten S, Ohno M, Mattaj JW: Nucleocytoplasmic transport: Ran, beta and beyond. *Trends Cell Biol* 2001, **11**:497-503.
48. Kudo N, Wolff B, Sekimoto T, Schreiner EP, Yoneda Y, Yanagida M, Horinouchi S, Yoshida M: Leptomycin B inhibition of signal-mediated nuclear export by direct binding to CRM1. *Exp Cell Res* 1998, **242**:540-547.
49. Mohr D, Frey S, Fischer T, Guttler T, Gorlich D: Characterisation of the passive permeability barrier of nuclear pore complexes. *EMBO J* 2009, **28**:2541-2553.
50. Stewart M, Baker RP, Bayliss R, Clayton L, Grant RP, Littlewood T, Matsuura Y: Molecular mechanism of translocation through nuclear pore complexes during nuclear protein import. *FEBS Lett* 2001, **498**:145-149.
51. Dabauvalle MC, Franke WW: Determination of the intracellular state of soluble macromolecules by gel filtration in vivo in the cytoplasm of amphibian oocytes. *J Cell Biol* 1986, **102**:2006-2014.
52. Hiebert SW, Lamb RA: Cell surface expression of glycosylated, nonglycosylated, and truncated forms of a cytoplasmic protein pyruvate kinase. *J Cell Biol* 1988, **107**:865-876.
53. Brameier M, Krings A, MacCallum RM: NucPred—predicting nuclear localization of proteins. *Bioinformatics* 2007, **23**:1159-1160.
54. Nair R, Rost B: LOC3D: annotate sub-cellular localization for protein structures. *Nucleic Acids Res* 2003, **31**:3337-3340.
55. la Cour T, Kierner L, Molgaard A, Gupta R, Skriver K, Brunak S: Analysis and prediction of leucine-rich nuclear export signals. *Protein Eng Des Sel* 2004, **17**:527-536.
56. Campbell EM, Perez O, Melar M, Hope TJ: Labeling HIV-1 virions with two fluorescent proteins allows identification of virions that have productively entered the target cell. *Virology* 2007, **360**:286-293.
57. Diaz-Griffero F, Li X, Javanbakht H, Song B, Welikala S, Stremmler M, Sodroski J: Rapid turnover and polyubiquitylation of the retroviral restriction factor TRIM5. *Virology* 2006, **349**:300-315.
58. Song B, Diaz-Griffero F, Park DH, Rogers T, Stremmler M, Sodroski J: TRIM5alpha association with cytoplasmic bodies is not required for antiretroviral activity. *Virology* 2005, **343**:201-211.
59. Reszka N, Zhou C, Song B, Sodroski JG, Knipe DM: Simian TRIM5alpha proteins reduce replication of herpes simplex virus. *Virology* 2010, **398**:243-250.
60. Diaz-Griffero F, Hoschander SA, Brojatsch J: Endocytosis is a critical step in entry of subgroup B avian leukosis viruses. *J Virol* 2002, **76**:12866-12876.

doi:10.1186/1742-4690-8-38

Cite this article as: Diaz-Griffero *et al.*: Trafficking of some old world primate TRIM5α proteins through the nucleus. *Retrovirology* 2011 **8**:38.