

MEETING ABSTRACT

Open Access

Stromal cells help ATL cells to escape from apoptosis induced by histone deacetylase inhibitors

Yukiko Miyatake^{1,2*}, Utano Tomaru¹, André L A Oliveira², Noreen Sheehy², Takashi Yoshiki¹, Masanori Kasahara¹, William W Hall²

From 15th International Conference on Human Retroviruses: HTLV and Related Viruses Leuven and Gembloux, Belgium. 5-8 June 2011

HTLV-1 expression is generally extremely low in freshly isolated ATL cells, suggesting that gene silencing has occurred. The silencing of viral genes in ATL cells could be due to not only host epigenetic mechanism, but also consequences of natural selection of ATL cells to evade the host immune response. Recently, it has been reported that stromal cells could contribute to the regulation of HTLV-1 expression. In this study, we investigated whether stromal cells could be involved in reactivation of viral gene expression in HTLV-1 infected cells using coculture systems. We used CR cells, a viral gene silenced ATL cell line, and human HEK-293T epithelial cells as stromal cells. To reactivate viral gene expression in CR cells, the histone deacetylase inhibitor, Trichostatin A (TSA) was employed. CR cells co-cultured with HEK-293T cells dramatically escaped TSA-induced apoptosis, while CR cells cultured alone underwent apoptosis by TSA in a dose-dependent manner. Moreover, the transcriptional activity of NF- κ B in CR cells co-cultured with HEK-293T cells was decreased by TSA, whereas the activity in CR cells cultured alone was increased. These results suggest that stromal cells might allow HTLV-1infected cells to escape from apoptosis induced by activation of HTLV-1 expression. This study provides an insight into the potential role of stromal cells in the maintenance of latency in HTLV-1 infected cells.

Author details

¹Department of Pathology, Hokkaido University Graduate School of Medicine, Sapporo, 060-8638, Japan. ²Centre for Research in Infectious

* Correspondence: yukimiya@med.hokudai.ac.jp

¹Department of Pathology, Hokkaido University Graduate School of Medicine, Sapporo, 060-8638, Japan
Full list of author information is available at the end of the article

Diseases, School of Medicine and Medical Science, University College Dublin, Dublin, 4, Ireland.

Published: 6 June 2011

doi:10.1186/1742-4690-8-S1-A167

Cite this article as: Miyatake *et al.*: Stromal cells help ATL cells to escape from apoptosis induced by histone deacetylase inhibitors. *Retrovirology* 2011 **8**(Suppl 1):A167.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at www.biomedcentral.com/submit



