



ORAL PRESENTATION

Open Access

Distinct gene expression profiles associated with the susceptibility of pathogen-specific CD4⁺ T cells to HIV-1 infection

H Hu^{1*}, M Nau¹, P Ehrenberg¹, A Chenine¹, Z Daye², Z Wei³, N Michael¹, M Vahey¹, J Kim¹, M Marovich¹, S Ratto-Kim¹

From AIDS Vaccine 2012
Boston, MA, USA. 9-12 September 2012

Background

HIV infection causes the progressive depletion of CD4⁺ T cells. Contrary to the early loss of CD4 response to opportunistic pathogens like *Candida albicans*, cytomegalovirus (CMV)-specific CD4 response is persistent when total CD4⁺ T cell number is low. The mechanism is less clear. Despite considerable knowledge for the impact of HIV infection on total CD4⁺ T cells and their subsets, little is known about HIV infection of CD4⁺ T cells of different pathogen/antigen (Ag) specificity.

Methods

PBMC from HIV-negative donors were CFSE-labeled and stimulated ex vivo with pathogen-specific antigens including viral (CMV), bacterial (Tetanus Toxoid: TT) and fungal (*Candida albicans*) antigens. HIV infection of Ag-specific CD4⁺ T cells was determined by intracellular p24 production in CFSE-low population.

Results

While TT- and *Candida*-specific CD4⁺ T cells were permissive, CMV-specific CD4⁺ T cells are highly resistant to both X4 and R5 HIV independent of coreceptor usage. Quantification of HIV DNA in sorted, antigen-specific CD4⁺ T cells demonstrated a reduction of both strong-stop and full-length HIV DNA in CMV-specific CD4⁺ T cells. β -chemokine neutralization enhanced HIV entry and viral replication in TT- and *Candida*-specific CD4⁺ T cells, whereas HIV infection in CMV-specific CD4⁺ T cells remained low despite increased HIV entry by β -chemokine neutralization, suggesting

both entry and post-entry HIV restriction in CMV-specific cells. Microarray analysis revealed distinct gene expression profiles that involved selective upregulation of a broad array of antiviral genes in CMV-specific CD4⁺ T cells, whereas TT- and *Candida*-specific CD4⁺ T cells mainly upregulated a Th17 inflammatory response.

Conclusion

Our data suggest a mechanism for the persistence of CMV-specific CD4 response and the earlier loss of mucosal Th17-associated TT- and *Candida*-specific CD4 response in AIDS patients. The model described is useful in HIV vaccine studies by evaluating the susceptibility of vaccine-specific CD4 responses to HIV infection.

Author details

¹U.S. Military HIV Research Program, Silver Spring, MD, USA. ²University of Pennsylvania School of Medicine, Philadelphia, PA, USA. ³New Jersey Institute of Technology, Newark, NJ, USA.

Published: 13 September 2012

doi:10.1186/1742-4690-9-S2-O46

Cite this article as: Hu et al.: Distinct gene expression profiles associated with the susceptibility of pathogen-specific CD4⁺ T cells to HIV-1 infection. *Retrovirology* 2012 **9**(Suppl 2):O46.

¹U.S. Military HIV Research Program, Silver Spring, MD, USA
Full list of author information is available at the end of the article